

ARUNACHAL Factsheet

Key Indicators

| SL No | A) Demographic Indicators (Census 2011) | | SL No | C) State Health Indicator (SRS 2014) | | | |
|---|--|------------------------|------------------------|--------------------------------------|--------------------------|----------------------------------|------|
| 1 | Population 2011 | 1,382,611 | 9 | Birth Rate | | | 19.3 |
| 2 | Sex Ratio | 920 | 10 | Death Rate | | | 5.8 |
| 3 | Number of districts | 16 | 11 | IMR | | | 32 |
| B) Infrastructure Details (RHS 2015) | | | 12 | Growth Rate | | | 13.4 |
| 4 | Sub Centers (SC) | 286 | 13 | TFR (SRS - 2007) | | | 2.7 |
| 5 | Primary Health Centers (PHC) | 117 | | | | | |
| 6 | Community Health Centers (CHC) | 52 | | | | | |
| 7 | SDH | 0 | | | | | |
| 8 | DH | 14 | | | | | |
| D) Maternal Health: | | 2015-16 | 2016-17 | | | | |
| 14 | Expected number of PW | 32,543 | 33,214 | | | | |
| 15 | Expected number of Delivery | 30,177 | 30,798 | | | | |
| | | HMIS (2015-16) | HMIS (2016-17) | | DLHS-4 (2012-13) | | |
| | | Number | PC | Number | PC | | |
| 16 | PW register for ANC | 29,631 | 91.1 | 29,946 | 90.2 | Any ANC | 61.6 |
| 17 | Mothers registered in the first trimester | 11,049 | 37.3 | 11,114 | 37.1 | ANC Check up in first trimester | 34.8 |
| 18 | Mothers who had at least 3 Ante-Natal care check ups | 12,771 | 43.1 | 12,414 | 41.5 | 3 or more ANC check up | 43.8 |
| 19 | PW who got TT 2 / Booster injection | 14,149 | 47.8 | 14,544 | 48.6 | Atleast 1 TT received | 59.1 |
| 20 | PW who consumed 100 IFA tablets | 18,866 | 63.7 | 30,671 | 102.4 | 100 IFA tablets consumed | 17.4 |
| 21 | Home Delivery against expected delivery * | 1,207 | 4.0 | 1,164 | 3.8 | Full ANC | 13.5 |
| 22 | Institutional Delivery against expected delivery * | 15,368 | 51 | 15,436 | 50 | Home Delivery | 48.5 |
| 23 | Home Delivery against reported delivery ** | 1,207 | 7 | 1,164 | 7.0 | Institutional Delivery | 49.5 |
| 24 | Institutional Delivery against reported delivery ** | 15,368 | 93 | 15,436 | 93.0 | Home Delivery by SBA | 3.1 |
| 25 | Total reported Deliveries | 16,575 | 55 | 16,600 | 54 | PNC within 14 days of delivery | 48.8 |
| 26 | Reported Delivery at home assisted by a Doctor/Nurse /LHV/ANM(%) | 692 | 57 | 456 | 39 | | |
| 27 | Woman getting post partum check up between 48 hours and 14 days to total reported deliveries | 4,997 | 30 | 4,488 | 27 | | |
| Note: In case of HMIS data, Percentage is with reference to total number of Pregnant Women registered for ANC during the period and in DLHS-4 is with ref. to Women who had live/still births since three years preceding the survey. (DLHS-4 reference period is from 1-1-2008 to survey date). | | | | | | | |
| 28 | Expected number of Live Birth | 29,585 | 30,195 | | | | |
| E) Child Immunization : | | HMIS (2015-16) | HMIS (2016-17) | | DLHS-4 (2012-13) | | |
| | | Number | PC | Number | PC | | |
| 29 | Total reported live births | 16,566 | 56.0 | 17,069 | 56.5 | Breastfed within 1 hour of birth | 55.7 |

| | | | | | | | |
|---|--|-----------------------|-----------------------|---------------------------|------|-------------------|------|
| 30 | Total Still Birth | 276 | 0.9 | 268 | 0.9 | BCG | 75.5 |
| 31 | Newborn weighted at birth to live birth | 15,189 | 91.7 | 16,261 | 95.3 | 3 dose of DPT | 71.1 |
| 32 | Newborn having weighted less than 2.5 kg to newborn weighted at birth. | 1,000 | 6.0 | 1,338 | 7.8 | Measles | 60.4 |
| 33 | Newborns breastfed within 1 hour to total reported live birth. | 14,551 | 87.8 | 13,920 | 81.6 | Full Immunization | 49.2 |
| 34 | Fully immunized | 17,052 | 57.6 | 18,131 | 60.0 | | |
| 35 | Children who have received BCG | 21,657 | 73.2 | 19,803 | 65.6 | | |
| 36 | Children who have received DPT1 | 17,680 | 59.8 | 1,320 | 4.4 | | |
| 37 | Children who have received DPT3 | 18,503 | 62.5 | 1,488 | 4.9 | | |
| 38 | Children who have received Measles | 18,750 | 63.4 | 17,925 | 59.4 | | |
| 39 | Diarrhoea and Dehydration cases reported in children below 5 years | 17,149 | | 15,830 | | | |
| Note: In case of HMIS data, Proportions are calculated for 0-11 months old infants to total number of reported live births and in DLHS-4, the percentage of children 12-23 months olds who received specific vaccination to mothers who had last and last but one living child born three years preceding the survey. (DLHS-4 reference period is from 1-1-2008 to survey date). | | | | | | | |
| F) Family Planning | | HMIS (2015-16) | HMIS (2016-17) | DLHS-4 (2012-13) | | | |
| 40 | Number of Vasectomies performed | 4 | 33 | Limiting | | 11.7 | |
| 41 | Number of Tubectomies performed | 968 | 1,069 | Spacing | | 20.4 | |
| 42 | Number of IUD insertion | 2,924 | 2,916 | Women using any method | | 48.4 | |
| 43 | Number of IUD removals | 1,333 | 1,627 | Women using modern method | | 43.9 | |
| G) Service delivery | | HMIS (2015-16) | HMIS (2016-17) | | | | |
| 44 | IPD cases per 10,000 population *** | 461 | 423 | | | | |
| 45 | General OPD cases per 10,000 population *** | 8,506 | 8733 | | | | |
| 46 | Ayush OPD cases per 10,000 population *** | 427 | 490 | | | | |
| H) Mortality Details | | | | | | | |
| I) | Infant Deaths (In Numbers) | HMIS (2015-16) | HMIS (2016-17) | | | | |
| 47 | Neonatal death within 24 hours | 5 | 14 | | | | |
| 48 | Up to 1 Weeks of Birth | 32 | 25 | | | | |
| 49 | Between 1 week & 4 weeks of birth | 0 | 0 | | | | |
| 50 | Neonatal Death | 37 | 39 | | | | |
| 51 | Between 1 month and 11 months | 29 | 29 | | | | |
| 52 | Total reported infant deaths | 66 | 68 | | | | |
| II) | Maternal Mortality | | | | | | |
| | Maternal Deaths (In Numbers) | HMIS (2015-16) | HMIS (2016-17) | | | | |
| 53 | Due to Abortion | 0 | 0 | | | | |
| 54 | Due to Obstructed/prolonged labour | 2 | 0 | | | | |
| 55 | Due to Severe hypertension/fits | 2 | 2 | | | | |
| 56 | Due to Bleeding | 3 | 0 | | | | |
| 57 | Due to High fever | 1 | 0 | | | | |
| 58 | Due to Other Causes (including causes not known) | 2 | 4 | | | | |
| 59 | Total reported maternal deaths | 10 | 6 | | | | |

Anjaw Factsheet

Key Indicators

| A) Demographic Indicators (Census 2011) | | | | | | | |
|--|--|----------------|------|----------------|-------|-----------------------------------|------|
| 1 | Population 2011 | 21,089 | | | | | |
| 2 | Sex Ratio | 805 | | | | | |
| B) Infrastructure Details (RHS 2015) | | Numbers | | | | | |
| 3 | Sub Centers (SC) | NA | | | | | |
| 4 | Primary Health Centers (PHC) | NA | | | | | |
| 5 | Community Health Centers (CHC) | 2 | | | | | |
| 6 | SDH | 0 | | | | | |
| 7 | DH | 0 | | | | | |
| C) Maternal Health: | | 2015-16 | | 2016-17 | | | |
| 8 | Expected number of PW | 475 | | 479 | | | |
| 9 | Expected number of Delivery | 440 | | 444 | | | |
| | | HMIS (2015-16) | | HMIS (2016-17) | | DLHS-4 (2012-13) | |
| | | Number | PC | Number | PC | | |
| 10 | PW register for ANC | 378 | 79.6 | 420 | 87.7 | Any ANC | 52.1 |
| 11 | Mothers registered in the first trimester | 135 | 35.7 | 159 | 37.9 | ANC Check up in first trimester | 22.1 |
| 12 | Mothers who had at least 3 Ante-Natal care check ups | 103 | 27.2 | 96 | 22.9 | 3 or more ANC check up | 32.6 |
| 13 | PW who got TT 2 / Booster injection | 154 | 40.7 | 172 | 41.0 | At least 1 TT received | 47.9 |
| 14 | PW who consumed 100 IFA tablets | 182 | 48.1 | 153 | 36.4 | 100 IFA tablets consumed | 10.3 |
| 15 | Home Delivery against expected delivery * | 19 | 4.3 | 0 | 0.0 | Full ANC | 6.5 |
| 16 | Institutional Delivery against expected delivery * | 102 | 23.2 | 94 | 21.2 | Home Delivery | 62.0 |
| 17 | Home Delivery against reported delivery ** | 19 | 15.7 | 0 | 0.0 | Institutional Delivery | 35.0 |
| 18 | Institutional Delivery against reported delivery ** | 102 | 84.3 | 94 | 100.0 | Home Delivery by SBA | 4.9 |
| 19 | Total reported Deliveries | 121 | | 94 | | PNC within 14 days of delivery | 65.2 |
| 20 | Reported Delivery at home assisted by a Doctor/Nurse /LHV/ANM(%) | 18 | 94.7 | 0 | - | | |
| 21 | Woman getting post partum check up between 48 hours and 14 days to total reported deliveries | 66 | 54.5 | 56 | 59.6 | | |
| Note: In case of HMIS data, Percentage is with reference to total number of Pregnant Women registered for ANC during the period and in DLHS-4 is with ref. to Women who had live/still births since three years preceding the survey. (DLHS-4 reference period is from 1-1-2008 to survey date). | | | | | | | |
| 22 | Expected number of Live Birth | 431 | | 435 | | | |
| D) Child Immunization : | | HMIS (2015-16) | | HMIS (2016-17) | | DLHS-4 (2012-13) | |
| 23 | Total reported live births | 119 | 27.6 | 94 | 21.6 | Breastfeed within 1 hour of birth | 53.8 |
| 24 | Total Still Birth | 2 | 0.5 | 0 | 0.0 | BCG | 65.5 |

| | | | | | | | |
|---|--|------------------------|------------------------|---------------------------|------|--------------------------|------|
| 25 | Newborn weighted at birth to live birth | 112 | 94.1 | 92 | 97.9 | 3 dose of DPT | 58.2 |
| 26 | Newborn having weight less than 2.5 kg to newborn weighted at birth. | 6 | 5.4 | 6 | 6.5 | Measles | 34.5 |
| 27 | Newborns breastfed within 1 hour to total reported live birth. | 111 | 93.3 | 93 | 98.9 | Full Immunization | 29.1 |
| 28 | Fully immunized | 131 | 30.4 | 104 | 23.9 | | |
| 29 | Children who have received BCG | 206 | 47.7 | 125 | 28.7 | | |
| 30 | Children who have received DPT1 | 256 | 59.3 | 9 | 2.1 | | |
| 31 | Children who have received DPT3 | 230 | 53.3 | 2 | 0.5 | | |
| 32 | Children who have received Measles | 252 | 58.4 | 125 | 28.7 | | |
| 33 | Diarrhoea and Dehydration cases reported in children below 5 years | 655 | | 626 | | | |
| <p>Note: In case of HMIS data, Proportions are calculated for 0-11 months old infants to total number of reported live births and in DLHS-4, the percentage of children 12-23 months olds who received specific vaccination to mothers who had last and last but one living child born three years preceding the survey. (DLHS-4 reference period is from 1-1-2008 to survey date).</p> | | | | | | | |
| E) Family Planning | | HMIS (2015-16) | HMIS (2016-17) | | | DLHS-4 (2012-13) | |
| 34 | Number of Vasectomies performed | 0 | 0 | Limiting | | 10.6 | |
| 35 | Number of Tubectomies performed | 0 | 0 | Spacing | | 27.1 | |
| 36 | Number of IUD insertion | 11 | 8 | Women using any method | | 44.4 | |
| 37 | Number of IUD removals | 6 | 8 | Women using modern method | | 39.1 | |
| F) Service delivery | | HMIS (2015-16) | HMIS (2016-17) | | | | |
| 38 | IPD cases per 10,000 population *** | 148 | 171 | | | | |
| 39 | General OPD cases per 10,000 population *** | 11479 | 11648 | | | | |
| 40 | Ayush OPD cases per 10,000 population *** | 875 | 905 | | | | |
| G) Mortality Details | | | | | | | |
| I) | Infant Deaths (In Numbers) | HMIS (2015-16) | HMIS (2016-17) | | | | |
| 41 | Neonatal death within 24 hours | 0 | 0 | | | | |
| 42 | Up to 1 Weeks of Birth | 0 | 0 | | | | |
| 43 | Between 1 week & 4 weeks of birth | 0 | 0 | | | | |
| 44 | Neonatal Death | 0 | 0 | | | | |
| 45 | Between 1 month and 11 months | 0 | 0 | | | | |
| 46 | Total reported infant deaths | 0 | 0 | | | | |
| II) | Maternal Mortality | | | | | | |
| | Maternal Deaths (In Numbers) | HMIS (2015-16) | HMIS (2016-17) | | | | |
| 47 | Due to Abortion | 0 | 0 | | | | |
| 48 | Due to Obstructed/prolonged labour | 0 | 0 | | | | |
| 49 | Due to Severe hypertension/fits | 0 | 0 | | | | |
| 50 | Due to Bleeding | 0 | 0 | | | | |
| 51 | Due to High fever | 0 | 0 | | | | |
| 52 | Due to Other Causes (including causes not known) | 0 | 0 | | | | |
| 53 | Total reported maternal deaths | 0 | 0 | | | | |

Changlang Factsheet

Key Indicators

| A) Demographic Indicators (Census 2011) | | | | | | | |
|--|--|-----------------|------|-----------------|------|-----------------------------------|------|
| 1 | Population 2011 | 147,951 | | | | | |
| 2 | Sex Ratio | 914 | | | | | |
| B) Infrastructure Details (RHS 2015) | | Numbers | | | | | |
| 3 | Sub Centers (SC) | NA | | | | | |
| 4 | Primary Health Centers (PHC) | NA | | | | | |
| 5 | Community Health Centers (CHC) | 4 | | | | | |
| 6 | SDH | 0 | | | | | |
| 7 | DH | 1 | | | | | |
| C) Maternal Health: | | 2015-16 | | 2016-17 | | | |
| 8 | Expected number of PW | 3,379 | | 3,422 | | | |
| 9 | Expected number of Delivery | 3,133 | | 3,173 | | | |
| | | HMIS (2015-16) | | HMIS (2016-17) | | DLHS-4 (2012-13) | |
| | | Number | PC | Number | PC | | |
| 10 | PW register for ANC | 3,219 | 95.3 | 3,154 | 92.2 | Any ANC | 63.2 |
| 11 | Mothers registered in the first trimester | 998 | 31.0 | 1,036 | 32.8 | ANC Check up in first trimester | 34.3 |
| 12 | Mothers who had at least 3 Ante-Natal care check ups | 1,470 | 45.7 | 1,566 | 49.7 | 3 or more ANC check up | 38.5 |
| 13 | PW who got TT 2 / Booster injection | 2,125 | 66.0 | 2,438 | 77.3 | At least 1 TT received | 60.5 |
| 14 | PW who consumed 100 IFA tablets | 1,331 | 41.3 | 1,610 | 51.0 | 100 IFA tablets consumed | 7.1 |
| 15 | Home Delivery against expected delivery * | 162 | 5.2 | 225 | 7.1 | Full ANC | 4.6 |
| 16 | Institutional Delivery against expected delivery * | 1,173 | 37.4 | 1,051 | 33.1 | Home Delivery | 46.7 |
| 17 | Home Delivery against reported delivery ** | 162 | 12.1 | 225 | 17.6 | Institutional Delivery | 52.2 |
| 18 | Institutional Delivery against reported delivery ** | 1,173 | 87.9 | 1,051 | 82.4 | Home Delivery by SBA | 0.0 |
| 19 | Total reported Deliveries | 1,335 | | 1,276 | | PNC within 14 days of delivery | 45.0 |
| 20 | Reported Delivery at home assisted by a Doctor/Nurse /LHV/ANM(%) | 121 | 74.7 | 84 | 37.3 | | |
| 21 | Woman getting post partum check up between 48 hours and 14 days to total reported deliveries | 550 | 41.2 | 452 | 35.4 | | |
| Note: In case of HMIS data, Percentage is with reference to total number of Pregnant Women registered for ANC during the period and in DLHS-4 is with ref. to Women who had live/still births since three years preceding the survey. (DLHS-4 reference period is from 1-1-2008 to survey date). | | | | | | | |
| 22 | Expected number of Live Birth | 3,072 | | 3,111 | | | |
| D) Child Immunization : | | HMIS (2015-16) | | HMIS (2016-17) | | DLHS-4 (2012-13) | |
| 23 | Total reported live births | 1,333 | 43.4 | 1,235 | 39.7 | Breastfeed within 1 hour of birth | 59.4 |
| 24 | Total Still Birth | 16 | 0.5 | 15 | 0.5 | BCG | 65.8 |

| | | | | | | | |
|---|--|-----------------------|-----------------------|---------------------------|------|-------------------------|------|
| 25 | Newborn weighted at birth to live birth | 1,262 | 94.7 | 1,183 | 95.8 | 3 dose of DPT | 72.6 |
| 26 | Newborn having weight less than 2.5 kg to newborn weighted at birth. | 113 | 9.0 | 83 | 7.0 | Measles | 47.9 |
| 27 | Newborns breastfed within 1 hour to total reported live birth. | 1,265 | 94.9 | 1,188 | 96.2 | Full Immunization | 36.1 |
| 28 | Fully immunized | 2,041 | 66.4 | 1,893 | 60.9 | | |
| 29 | Children who have received BCG | 2,630 | 85.6 | 2,302 | 74.0 | | |
| 30 | Children who have received DPT1 | 2,220 | 72.3 | 72 | 2.3 | | |
| 31 | Children who have received DPT3 | 2,249 | 73.2 | 197 | 6.3 | | |
| 32 | Children who have received Measles | 2,367 | 77.1 | 2,097 | 67.4 | | |
| 33 | Diarrhoea and Dehydration cases reported in children below 5 years | 364 | | 386 | | | |
| <p>Note: In case of HMIS data, Proportions are calculated for 0-11 months old infants to total number of reported live births and in DLHS-4, the percentage of children 12-23 months olds who received specific vaccination to mothers who had last and last but one living child born three years preceding the survey. (DLHS-4 reference period is from 1-1-2008 to survey date).</p> | | | | | | | |
| E) Family Planning | | HMIS (2015-16) | HMIS (2016-17) | | | DLHS-4 (2012-13) | |
| 34 | Number of Vasectomies performed | 0 | 0 | Limiting | | 13.2 | |
| 35 | Number of Tubectomies performed | 49 | 0 | Spacing | | 24 | |
| 36 | Number of IUD insertion | 104 | 125 | Women using any method | | 44.7 | |
| 37 | Number of IUD removals | 51 | 145 | Women using modern method | | 38.5 | |
| F) Service delivery | | HMIS (2015-16) | HMIS (2016-17) | | | | |
| 38 | IPD cases per 10,000 population *** | 196 | 226 | | | | |
| 39 | General OPD cases per 10,000 population *** | 6259 | 7306 | | | | |
| 40 | Ayush OPD cases per 10,000 population *** | 742 | 725 | | | | |
| G) Mortality Details | | | | | | | |
| I) | Infant Deaths (In Numbers) | HMIS (2015-16) | HMIS (2016-17) | | | | |
| 41 | Neonatal death within 24 hours | 0 | 0 | | | | |
| 42 | Up to 1 Weeks of Birth | 1 | 0 | | | | |
| 43 | Between 1 week & 4 weeks of birth | 0 | 0 | | | | |
| 44 | Neonatal Death | 1 | 0 | | | | |
| 45 | Between 1 month and 11 months | 0 | 0 | | | | |
| 46 | Total reported infant deaths | 1 | 0 | | | | |
| II) | Maternal Mortality | | | | | | |
| | Maternal Deaths (In Numbers) | HMIS (2015-16) | HMIS (2016-17) | | | | |
| 47 | Due to Abortion | 0 | 0 | | | | |
| 48 | Due to Obstructed/prolonged labour | 0 | 0 | | | | |
| 49 | Due to Severe hypertension/fits | 0 | 0 | | | | |
| 50 | Due to Bleeding | 0 | 0 | | | | |
| 51 | Due to High fever | 0 | 0 | | | | |
| 52 | Due to Other Causes (including causes not known) | 0 | 0 | | | | |
| 53 | Total reported maternal deaths | 0 | 0 | | | | |

Dibang Valley Factsheet

Key Indicators

| A) Demographic Indicators (Census 2011) | | | | | | | |
|--|--|----------------|-------|----------------|-------|-----------------------------------|------|
| 1 | Population 2011 | 7,948 | | | | | |
| 2 | Sex Ratio | 808 | | | | | |
| B) Infrastructure Details (RHS 2015) | | Numbers | | | | | |
| 3 | Sub Centers (SC) | NA | | | | | |
| 4 | Primary Health Centers (PHC) | NA | | | | | |
| 5 | Community Health Centers (CHC) | 0 | | | | | |
| 6 | SDH | 0 | | | | | |
| 7 | DH | 1 | | | | | |
| C) Maternal Health: | | 2015-16 | | 2016-17 | | | |
| 8 | Expected number of PW | 176 | | 177 | | | |
| 9 | Expected number of Delivery | 164 | | 164 | | | |
| | | HMIS (2015-16) | | HMIS (2016-17) | | DLHS-4 (2012-13) | |
| | | Number | PC | Number | PC | | |
| 10 | PW register for ANC | 175 | 99.2 | 200 | 113.0 | Any ANC | 45.6 |
| 11 | Mothers registered in the first trimester | 103 | 58.9 | 115 | 57.5 | ANC Check up in first trimester | 25.0 |
| 12 | Mothers who had at least 3 Ante-Natal care check ups | 72 | 41.1 | 52 | 26.0 | 3 or more ANC check up | 31.7 |
| 13 | PW who got TT 2 / Booster injection | 90 | 51.4 | 104 | 52.0 | At least 1 TT received | 45.6 |
| 14 | PW who consumed 100 IFA tablets | 152 | 86.9 | 137 | 68.5 | 100 IFA tablets consumed | 12.8 |
| 15 | Home Delivery against expected delivery * | 3 | 1.8 | 9 | 5.5 | Full ANC | 9.5 |
| 16 | Institutional Delivery against expected delivery * | 73 | 44.6 | 61 | 37.2 | Home Delivery | 47.2 |
| 17 | Home Delivery against reported delivery ** | 3 | 3.9 | 9 | 12.9 | Institutional Delivery | 52.5 |
| 18 | Institutional Delivery against reported delivery ** | 73 | 96.1 | 61 | 87.1 | Home Delivery by SBA | 3.4 |
| 19 | Total reported Deliveries | 76 | | 70 | | PNC within 14 days of delivery | 55.3 |
| 20 | Reported Delivery at home assisted by a Doctor/Nurse /LHV/ANM(%) | 3 | 100.0 | 7 | 77.8 | | |
| 21 | Woman getting post partum check up between 48 hours and 14 days to total reported deliveries | 72 | 94.7 | 67 | 95.7 | | |
| Note: In case of HMIS data, Percentage is with reference to total number of Pregnant Women registered for ANC during the period and in DLHS-4 is with ref. to Women who had live/still births since three years preceding the survey. (DLHS-4 reference period is from 1-1-2008 to survey date). | | | | | | | |
| 22 | Expected number of Live Birth | 160 | | 161 | | | |
| D) Child Immunization : | | HMIS (2015-16) | | HMIS (2016-17) | | DLHS-4 (2012-13) | |
| 23 | Total reported live births | 80 | 49.9 | 74 | 46.0 | Breastfeed within 1 hour of birth | 50.4 |
| 24 | Total Still Birth | 0 | 0.0 | 3 | 1.9 | BCG | 90.2 |

| | | | | | | | |
|---|--|------------------------|------|------------------------|------|---------------------------|------|
| 25 | Newborn weighted at birth to live birth | 79 | 98.8 | 71 | 95.9 | 3 dose of DPT | 90.2 |
| 26 | Newborn having weight less than 2.5 kg to newborn weighted at birth. | 2 | 2.5 | 4 | 5.6 | Measles | 61 |
| 27 | Newborns breastfed within 1 hour to total reported live birth. | 79 | 98.8 | 71 | 95.9 | Full Immunization | 58.5 |
| 28 | Fully immunized | 89 | 55.5 | 36 | 22.4 | | |
| 29 | Children who have received BCG | 81 | 50.5 | 69 | 42.9 | | |
| 30 | Children who have received DPT1 | 78 | 48.7 | 11 | 6.8 | | |
| 31 | Children who have received DPT3 | 81 | 50.5 | 9 | 5.6 | | |
| 32 | Children who have received Measles | 91 | 56.8 | 61 | 37.9 | | |
| 33 | Diarrhoea and Dehydration cases reported in children below 5 years | 375 | | 217 | | | |
| <p>Note: In case of HMIS data, Proportions are calculated for 0-11 months old infants to total number of reported live births and in DLHS-4, the percentage of children 12-23 months olds who received specific vaccination to mothers who had last and last but one living child born three years preceding the survey. (DLHS-4 reference period is from 1-1-2008 to survey date).</p> | | | | | | | |
| E) Family Planning | | HMIS (2015-16) | | HMIS (2016-17) | | DLHS-4 (2012-13) | |
| 34 | Number of Vasectomies performed | 0 | | 0 | | Limiting | 10.2 |
| 35 | Number of Tubectomies performed | 0 | | 0 | | Spacing | 24.5 |
| 36 | Number of IUD insertion | 1 | | 3 | | Women using any method | 45.4 |
| 37 | Number of IUD removals | 0 | | 3 | | Women using modern method | 44.3 |
| F) Service delivery | | HMIS (2015-16) | | HMIS (2016-17) | | | |
| 38 | IPD cases per 10,000 population *** | 258 | | 89 | | | |
| 39 | General OPD cases per 10,000 population *** | 13026 | | 11899 | | | |
| 40 | Ayush OPD cases per 10,000 population *** | 462 | | 1403 | | | |
| G) Mortality Details | | | | | | | |
| I) | Infant Deaths (In Numbers) | HMIS (2015-16) | | HMIS (2016-17) | | | |
| 41 | Neonatal death within 24 hours | 0 | | 0 | | | |
| 42 | Up to 1 Weeks of Birth | 0 | | 0 | | | |
| 43 | Between 1 week & 4 weeks of birth | 0 | | 0 | | | |
| 44 | Neonatal Death | 0 | | 0 | | | |
| 45 | Between 1 month and 11 months | 0 | | 0 | | | |
| 46 | Total reported infant deaths | 0 | | 0 | | | |
| II) | Maternal Mortality | HMIS (2015-16) | | HMIS (2016-17) | | | |
| | Maternal Deaths (In Numbers) | HMIS (2015-16) | | HMIS (2016-17) | | | |
| 47 | Due to Abortion | 0 | | 0 | | | |
| 48 | Due to Obstructed/prolonged labour | 0 | | 0 | | | |
| 49 | Due to Severe hypertension/fits | 0 | | 0 | | | |
| 50 | Due to Bleeding | 0 | | 0 | | | |
| 51 | Due to High fever | 0 | | 0 | | | |
| 52 | Due to Other Causes (including causes not known) | 0 | | 0 | | | |
| 53 | Total reported maternal deaths | 0 | | 0 | | | |

East Kameng Factsheet

Key Indicators

| A) Demographic Indicators (Census 2011) | | | | | | | |
|--|--|----------------|-------|----------------|------|-----------------------------------|------|
| 1 | Population 2011 | 78,413 | | | | | |
| 2 | Sex Ratio | 1,012 | | | | | |
| B) Infrastructure Details (RHS 2015) | | Numbers | | | | | |
| 3 | Sub Centers (SC) | NA | | | | | |
| 4 | Primary Health Centers (PHC) | NA | | | | | |
| 5 | Community Health Centers (CHC) | 3 | | | | | |
| 6 | SDH | 0 | | | | | |
| 7 | DH | 1 | | | | | |
| C) Maternal Health: | | 2015-16 | | 2016-17 | | | |
| 8 | Expected number of PW | 1,933 | | 1,994 | | | |
| 9 | Expected number of Delivery | 1,792 | | 1,849 | | | |
| | | HMIS (2015-16) | | HMIS (2016-17) | | DLHS-4 (2012-13) | |
| | | Number | PC | Number | PC | | |
| 10 | PW register for ANC | 2,179 | 112.7 | 1,712 | 85.8 | Any ANC | 48.0 |
| 11 | Mothers registered in the first trimester | 848 | 38.9 | 499 | 29.1 | ANC Check up in first trimester | 28.9 |
| 12 | Mothers who had at least 3 Ante-Natal care check ups | 822 | 37.7 | 780 | 45.6 | 3 or more ANC check up | 29.6 |
| 13 | PW who got TT 2 / Booster injection | 958 | 44.0 | 919 | 53.7 | At least 1 TT received | 44.4 |
| 14 | PW who consumed 100 IFA tablets | 1,215 | 55.8 | 400 | 23.4 | 100 IFA tablets consumed | 12.1 |
| 15 | Home Delivery against expected delivery * | 94 | 5.2 | 37 | 2.0 | Full ANC | 7.5 |
| 16 | Institutional Delivery against expected delivery * | 652 | 36.4 | 620 | 33.5 | Home Delivery | 59.5 |
| 17 | Home Delivery against reported delivery ** | 94 | 12.6 | 37 | 5.6 | Institutional Delivery | 34.1 |
| 18 | Institutional Delivery against reported delivery ** | 652 | 87.4 | 620 | 94.4 | Home Delivery by SBA | 2.2 |
| 19 | Total reported Deliveries | 746 | | 657 | | PNC within 14 days of delivery | 52.6 |
| 20 | Reported Delivery at home assisted by a Doctor/Nurse /LHV/ANM(%) | 74 | 78.7 | 23 | 62.2 | | |
| 21 | Woman getting post partum check up between 48 hours and 14 days to total reported deliveries | 71 | 9.5 | 68 | 10.4 | | |
| Note: In case of HMIS data, Percentage is with reference to total number of Pregnant Women registered for ANC during the period and in DLHS-4 is with ref. to Women who had live/still births since three years preceding the survey. (DLHS-4 reference period is from 1-1-2008 to survey date). | | | | | | | |
| 22 | Expected number of Live Birth | 1,757 | | 1,813 | | | |
| D) Child Immunization : | | HMIS (2015-16) | | HMIS (2016-17) | | DLHS-4 (2012-13) | |
| 23 | Total reported live births | 785 | 44.7 | 773 | 42.6 | Breastfeed within 1 hour of birth | 45.9 |
| 24 | Total Still Birth | 14 | 0.8 | 10 | 0.6 | BCG | 55.9 |

| | | | | | | | |
|---|--|------------------------|-------|------------------------|------|---------------------------|------|
| 25 | Newborn weighted at birth to live birth | 681 | 86.8 | 572 | 74.0 | 3 dose of DPT | 46.3 |
| 26 | Newborn having weight less than 2.5 kg to newborn weighted at birth. | 143 | 21.0 | 22 | 3.8 | Measles | 40.3 |
| 27 | Newborns breastfed within 1 hour to total reported live birth. | 405 | 51.6 | 532 | 68.8 | Full Immunization | 18.2 |
| 28 | Fully immunized | 962 | 54.7 | 797 | 44.0 | | |
| 29 | Children who have received BCG | 1,218 | 69.3 | 876 | 48.3 | | |
| 30 | Children who have received DPT1 | 1,617 | 92.0 | 72 | 4.0 | | |
| 31 | Children who have received DPT3 | 1,985 | 113.0 | 89 | 4.9 | | |
| 32 | Children who have received Measles | 1,278 | 72.7 | 731 | 40.3 | | |
| 33 | Diarrhoea and Dehydration cases reported in children below 5 years | 2,414 | | 1,748 | | | |
| <p>Note: In case of HMIS data, Proportions are calculated for 0-11 months old infants to total number of reported live births and in DLHS-4, the percentage of children 12-23 months olds who received specific vaccination to mothers who had last and last but one living child born three years preceding the survey. (DLHS-4 reference period is from 1-1-2008 to survey date).</p> | | | | | | | |
| E) Family Planning | | HMIS (2015-16) | | HMIS (2016-17) | | DLHS-4 (2012-13) | |
| 34 | Number of Vasectomies performed | 0 | | 0 | | Limiting | 10.4 |
| 35 | Number of Tubectomies performed | 107 | | 0 | | Spacing | 24 |
| 36 | Number of IUD insertion | 188 | | 198 | | Women using any method | 44.5 |
| 37 | Number of IUD removals | 40 | | 134 | | Women using modern method | 38.7 |
| F) Service delivery | | HMIS (2015-16) | | HMIS (2016-17) | | | |
| 38 | IPD cases per 10,000 population *** | 335 | | 349 | | | |
| 39 | General OPD cases per 10,000 population *** | 4988 | | 5386 | | | |
| 40 | Ayush OPD cases per 10,000 population *** | 154 | | 446 | | | |
| G) Mortality Details | | | | | | | |
| I) | Infant Deaths (In Numbers) | HMIS (2015-16) | | HMIS (2016-17) | | | |
| 41 | Neonatal death within 24 hours | 0 | | 0 | | | |
| 42 | Up to 1 Weeks of Birth | 2 | | 0 | | | |
| 43 | Between 1 week & 4 weeks of birth | 0 | | 0 | | | |
| 44 | Neonatal Death | 2 | | 0 | | | |
| 45 | Between 1 month and 11 months | 6 | | 6 | | | |
| 46 | Total reported infant deaths | 8 | | 6 | | | |
| II) | Maternal Mortality | HMIS (2015-16) | | HMIS (2016-17) | | | |
| | Maternal Deaths (In Numbers) | HMIS (2015-16) | | HMIS (2016-17) | | | |
| 47 | Due to Abortion | 0 | | 0 | | | |
| 48 | Due to Obstructed/prolonged labour | 0 | | 0 | | | |
| 49 | Due to Severe hypertension/fits | 0 | | 1 | | | |
| 50 | Due to Bleeding | 0 | | 0 | | | |
| 51 | Due to High fever | 0 | | 0 | | | |
| 52 | Due to Other Causes (including causes not known) | 0 | | 0 | | | |
| 53 | Total reported maternal deaths | 0 | | 1 | | | |

East Siang Factsheet

Key Indicators

| A) Demographic Indicators (Census 2011) | | | | | | | |
|--|--|-----------------|------|-----------------|------|-----------------------------------|------|
| 1 | Population 2011 | 99,019 | | | | | |
| 2 | Sex Ratio | 962 | | | | | |
| B) Infrastructure Details (RHS 2015) | | Numbers | | | | | |
| 3 | Sub Centers (SC) | NA | | | | | |
| 4 | Primary Health Centers (PHC) | NA | | | | | |
| 5 | Community Health Centers (CHC) | 5 | | | | | |
| 6 | SDH | 0 | | | | | |
| 7 | DH | 1 | | | | | |
| C) Maternal Health: | | 2015-16 | | 2016-17 | | | |
| 8 | Expected number of PW | 2,221 | | 2,238 | | | |
| 9 | Expected number of Delivery | 2,059 | | 2,076 | | | |
| | | HMIS (2015-16) | | HMIS (2016-17) | | DLHS-4 (2012-13) | |
| | | Number | PC | Number | PC | | |
| 10 | PW register for ANC | 1,848 | 83.2 | 1,932 | 86.3 | Any ANC | 63.1 |
| 11 | Mothers registered in the first trimester | 1,002 | 54.2 | 1,091 | 56.5 | ANC Check up in first trimester | 37.1 |
| 12 | Mothers who had at least 3 Ante-Natal care check ups | 926 | 50.1 | 949 | 49.1 | 3 or more ANC check up | 50.2 |
| 13 | PW who got TT 2 / Booster injection | 1,031 | 55.8 | 1,134 | 58.7 | At least 1 TT received | 57.9 |
| 14 | PW who consumed 100 IFA tablets | 995 | 53.8 | 1,004 | 52.0 | 100 IFA tablets consumed | 19.3 |
| 15 | Home Delivery against expected delivery * | 56 | 2.7 | 52 | 2.5 | Full ANC | 13.8 |
| 16 | Institutional Delivery against expected delivery * | 1,631 | 79.2 | 1,743 | 84.0 | Home Delivery | 48.9 |
| 17 | Home Delivery against reported delivery ** | 56 | 3.3 | 52 | 2.9 | Institutional Delivery | 50.2 |
| 18 | Institutional Delivery against reported delivery ** | 1,631 | 96.7 | 1,743 | 97.1 | Home Delivery by SBA | 2.1 |
| 19 | Total reported Deliveries | 1,687 | | 1,795 | | PNC within 14 days of delivery | 50.8 |
| 20 | Reported Delivery at home assisted by a Doctor/Nurse /LHV/ANM(%) | 39 | 69.6 | 51 | 98.1 | | |
| 21 | Woman getting post partum check up between 48 hours and 14 days to total reported deliveries | 358 | 21.2 | 236 | 13.1 | | |
| Note: In case of HMIS data, Percentage is with reference to total number of Pregnant Women registered for ANC during the period and in DLHS-4 is with ref. to Women who had live/still births since three years preceding the survey. (DLHS-4 reference period is from 1-1-2008 to survey date). | | | | | | | |
| 22 | Expected number of Live Birth | 2,019 | | 2,035 | | | |
| D) Child Immunization : | | HMIS (2015-16) | | HMIS (2016-17) | | DLHS-4 (2012-13) | |
| 23 | Total reported live births | 1,655 | 82.0 | 1,790 | 88.0 | Breastfeed within 1 hour of birth | 53.3 |
| 24 | Total Still Birth | 32 | 1.6 | 47 | 2.3 | BCG | 69.8 |

| | | | | | | | |
|---|--|------------------------|------|------------------------|------|---------------------------|------|
| 25 | Newborn weighted at birth to live birth | 1,544 | 93.3 | 1,763 | 98.5 | 3 dose of DPT | 73.6 |
| 26 | Newborn having weight less than 2.5 kg to newborn weighted at birth. | 101 | 6.5 | 266 | 15.1 | Measles | 60.4 |
| 27 | Newborns breastfed within 1 hour to total reported live birth. | 683 | 41.3 | 626 | 35.0 | Full Immunization | 41.5 |
| 28 | Fully immunized | 1,278 | 63.3 | 1,245 | 61.2 | | |
| 29 | Children who have received BCG | 1,974 | 97.8 | 1,870 | 91.9 | | |
| 30 | Children who have received DPT1 | 1,301 | 64.4 | 56 | 2.8 | | |
| 31 | Children who have received DPT3 | 1,435 | 71.1 | 51 | 2.5 | | |
| 32 | Children who have received Measles | 1,441 | 71.4 | 1,447 | 71.1 | | |
| 33 | Diarrhoea and Dehydration cases reported in children below 5 years | 991 | | 534 | | | |
| <p>Note: In case of HMIS data, Proportions are calculated for 0-11 months old infants to total number of reported live births and in DLHS-4, the percentage of children 12-23 months olds who received specific vaccination to mothers who had last and last but one living child born three years preceding the survey. (DLHS-4 reference period is from 1-1-2008 to survey date).</p> | | | | | | | |
| E) Family Planning | | HMIS (2015-16) | | HMIS (2016-17) | | DLHS-4 (2012-13) | |
| 34 | Number of Vasectomies performed | 2 | | 4 | | Limiting | 12.7 |
| 35 | Number of Tubectomies performed | 212 | | 264 | | Spacing | 18.3 |
| 36 | Number of IUD insertion | 177 | | 198 | | Women using any method | 55.3 |
| 37 | Number of IUD removals | 75 | | 96 | | Women using modern method | 50 |
| F) Service delivery | | HMIS (2015-16) | | HMIS (2016-17) | | | |
| 38 | IPD cases per 10,000 population *** | 1003 | | 967 | | | |
| 39 | General OPD cases per 10,000 population *** | 14452 | | 15554 | | | |
| 40 | Ayush OPD cases per 10,000 population *** | 178 | | 187 | | | |
| G) Mortality Details | | | | | | | |
| I) | Infant Deaths (In Numbers) | HMIS (2015-16) | | HMIS (2016-17) | | | |
| 41 | Neonatal death within 24 hours | 1 | | 4 | | | |
| 42 | Up to 1 Weeks of Birth | 7 | | 5 | | | |
| 43 | Between 1 week & 4 weeks of birth | 0 | | 0 | | | |
| 44 | Neonatal Death | 8 | | 9 | | | |
| 45 | Between 1 month and 11 months | 2 | | 11 | | | |
| 46 | Total reported infant deaths | 10 | | 20 | | | |
| II) | Maternal Mortality | | | | | | |
| | Maternal Deaths (In Numbers) | HMIS (2015-16) | | HMIS (2016-17) | | | |
| 47 | Due to Abortion | 0 | | 0 | | | |
| 48 | Due to Obstructed/prolonged labour | 0 | | 0 | | | |
| 49 | Due to Severe hypertension/fits | 0 | | 0 | | | |
| 50 | Due to Bleeding | 0 | | 0 | | | |
| 51 | Due to High fever | 0 | | 0 | | | |
| 52 | Due to Other Causes (including causes not known) | 0 | | 2 | | | |
| 53 | Total reported maternal deaths | 0 | | 2 | | | |

Kurung Kumey Factsheet

Key Indicators

| A) Demographic Indicators (Census 2011) | | | | | | | |
|--|--|-----------------|------|-----------------|------|-----------------------------------|------|
| 1 | Population 2011 | 89,717 | | | | | |
| 2 | Sex Ratio | 1,029 | | | | | |
| B) Infrastructure Details (RHS 2015) | | Numbers | | | | | |
| 3 | Sub Centers (SC) | NA | | | | | |
| 4 | Primary Health Centers (PHC) | NA | | | | | |
| 5 | Community Health Centers (CHC) | 3 | | | | | |
| 6 | SDH | 0 | | | | | |
| 7 | DH | 0 | | | | | |
| C) Maternal Health: | | 2015-16 | | 2016-17 | | | |
| 8 | Expected number of PW | 2,978 | | 3,292 | | | |
| 9 | Expected number of Delivery | 2,762 | | 3,052 | | | |
| | | HMIS (2015-16) | | HMIS (2016-17) | | DLHS-4 (2012-13) | |
| | | Number | PC | Number | PC | | |
| 10 | PW register for ANC | 1,181 | 39.7 | 1,047 | 31.8 | Any ANC | 46.9 |
| 11 | Mothers registered in the first trimester | 656 | 55.5 | 643 | 61.4 | ANC Check up in first trimester | 21.3 |
| 12 | Mothers who had at least 3 Ante-Natal care check ups | 623 | 52.8 | 593 | 56.6 | 3 or more ANC check up | 23.9 |
| 13 | PW who got TT 2 / Booster injection | 409 | 34.6 | 397 | 37.9 | At least 1 TT received | 44.4 |
| 14 | PW who consumed 100 IFA tablets | 1,162 | 98.4 | 1,020 | 97.4 | 100 IFA tablets consumed | 13.8 |
| 15 | Home Delivery against expected delivery * | 160 | 5.8 | 150 | 4.9 | Full ANC | 8.4 |
| 16 | Institutional Delivery against expected delivery * | 396 | 14.3 | 323 | 10.6 | Home Delivery | 66.4 |
| 17 | Home Delivery against reported delivery ** | 160 | 28.8 | 150 | 31.7 | Institutional Delivery | 31.7 |
| 18 | Institutional Delivery against reported delivery ** | 396 | 71.2 | 323 | 68.3 | Home Delivery by SBA | 4.2 |
| 19 | Total reported Deliveries | 556 | | 473 | | PNC within 14 days of delivery | 39.5 |
| 20 | Reported Delivery at home assisted by a Doctor/Nurse /LHV/ANM(%) | 95 | 59.4 | 70 | 46.7 | | |
| 21 | Woman getting post partum check up between 48 hours and 14 days to total reported deliveries | 342 | 61.5 | 281 | 59.4 | | |
| Note: In case of HMIS data, Percentage is with reference to total number of Pregnant Women registered for ANC during the period and in DLHS-4 is with ref. to Women who had live/still births since three years preceding the survey. (DLHS-4 reference period is from 1-1-2008 to survey date). | | | | | | | |
| 22 | Expected number of Live Birth | 2,708 | | 2,993 | | | |
| D) Child Immunization : | | HMIS (2015-16) | | HMIS (2016-17) | | DLHS-4 (2012-13) | |
| 23 | Total reported live births | 561 | 20.7 | 501 | 16.7 | Breastfeed within 1 hour of birth | 67.7 |
| 24 | Total Still Birth | 1 | 0.0 | 9 | 0.3 | BCG | 77.6 |

| | | | | | | | |
|---|--|------------------------|------|------------------------|------|---------------------------|------|
| 25 | Newborn weighted at birth to live birth | 554 | 98.8 | 468 | 93.4 | 3 dose of DPT | 76.1 |
| 26 | Newborn having weight less than 2.5 kg to newborn weighted at birth. | 1 | 0.2 | 16 | 3.4 | Measles | 59.7 |
| 27 | Newborns breastfed within 1 hour to total reported live birth. | 560 | 99.8 | 464 | 92.6 | Full Immunization | 52.2 |
| 28 | Fully immunized | 810 | 29.9 | 815 | 27.2 | | |
| 29 | Children who have received BCG | 994 | 36.7 | 794 | 26.5 | | |
| 30 | Children who have received DPT1 | 875 | 32.3 | 129 | 4.3 | | |
| 31 | Children who have received DPT3 | 841 | 31.1 | 124 | 4.1 | | |
| 32 | Children who have received Measles | 811 | 30.0 | 800 | 26.7 | | |
| 33 | Diarrhoea and Dehydration cases reported in children below 5 years | 1,030 | | 786 | | | |
| <p>Note: In case of HMIS data, Proportions are calculated for 0-11 months old infants to total number of reported live births and in DLHS-4, the percentage of children 12-23 months olds who received specific vaccination to mothers who had last and last but one living child born three years preceding the survey. (DLHS-4 reference period is from 1-1-2008 to survey date).</p> | | | | | | | |
| E) Family Planning | | HMIS (2015-16) | | HMIS (2016-17) | | DLHS-4 (2012-13) | |
| 34 | Number of Vasectomies performed | 0 | | 0 | | Limiting | 11.4 |
| 35 | Number of Tubectomies performed | 0 | | 0 | | Spacing | 29 |
| 36 | Number of IUD insertion | 99 | | 133 | | Women using any method | 34.1 |
| 37 | Number of IUD removals | 31 | | 77 | | Women using modern method | 29.7 |
| F) Service delivery | | HMIS (2015-16) | | HMIS (2016-17) | | | |
| 38 | IPD cases per 10,000 population *** | 139 | | 107 | | | |
| 39 | General OPD cases per 10,000 population *** | 2185 | | 1915 | | | |
| 40 | Ayush OPD cases per 10,000 population *** | 91 | | 31 | | | |
| G) Mortality Details | | | | | | | |
| I) | Infant Deaths (In Numbers) | HMIS (2015-16) | | HMIS (2016-17) | | | |
| 41 | Neonatal death within 24 hours | 0 | | 0 | | | |
| 42 | Up to 1 Weeks of Birth | 0 | | 0 | | | |
| 43 | Between 1 week & 4 weeks of birth | 0 | | 0 | | | |
| 44 | Neonatal Death | 0 | | 0 | | | |
| 45 | Between 1 month and 11 months | 0 | | 0 | | | |
| 46 | Total reported infant deaths | 0 | | 0 | | | |
| II) | Maternal Mortality | HMIS (2015-16) | | HMIS (2016-17) | | | |
| | Maternal Deaths (In Numbers) | HMIS (2015-16) | | HMIS (2016-17) | | | |
| 47 | Due to Abortion | 0 | | 0 | | | |
| 48 | Due to Obstructed/prolonged labour | 0 | | 0 | | | |
| 49 | Due to Severe hypertension/fits | 0 | | 0 | | | |
| 50 | Due to Bleeding | 0 | | 0 | | | |
| 51 | Due to High fever | 0 | | 0 | | | |
| 52 | Due to Other Causes (including causes not known) | 0 | | 0 | | | |
| 53 | Total reported maternal deaths | 0 | | 0 | | | |

Lohit Factsheet

Key Indicators

| A) Demographic Indicators (Census 2011) | | | | | | | |
|--|--|----------------|-------|----------------|-------|-----------------------------------|------|
| 1 | Population 2011 | 145,538 | | | | | |
| 2 | Sex Ratio | 901 | | | | | |
| B) Infrastructure Details (RHS 2015) | | Numbers | | | | | |
| 3 | Sub Centers (SC) | NA | | | | | |
| 4 | Primary Health Centers (PHC) | NA | | | | | |
| 5 | Community Health Centers (CHC) | 3 | | | | | |
| 6 | SDH | 0 | | | | | |
| 7 | DH | 1 | | | | | |
| C) Maternal Health: | | 2015-16 | | 2016-17 | | | |
| 8 | Expected number of PW | 3,302 | | 3,339 | | | |
| 9 | Expected number of Delivery | 3,062 | | 3,096 | | | |
| | | HMIS (2015-16) | | HMIS (2016-17) | | DLHS-4 (2012-13) | |
| | | Number | PC | Number | PC | | |
| 10 | PW register for ANC | 3,455 | 104.6 | 3,568 | 106.9 | Any ANC | 62.8 |
| 11 | Mothers registered in the first trimester | 1,319 | 38.2 | 1,363 | 38.2 | ANC Check up in first trimester | 32.5 |
| 12 | Mothers who had at least 3 Ante-Natal care check ups | 1,632 | 47.2 | 1,652 | 46.3 | 3 or more ANC check up | 45.8 |
| 13 | PW who got TT 2 / Booster injection | 2,136 | 61.8 | 2,162 | 60.6 | At least 1 TT received | 60.6 |
| 14 | PW who consumed 100 IFA tablets | 5,606 | 162.3 | 10,238 | 286.9 | 100 IFA tablets consumed | 15.8 |
| 15 | Home Delivery against expected delivery * | 19 | 0.6 | 33 | 1.1 | Full ANC | 12.0 |
| 16 | Institutional Delivery against expected delivery * | 1,543 | 50.4 | 1,521 | 49.1 | Home Delivery | 49.5 |
| 17 | Home Delivery against reported delivery ** | 19 | 1.2 | 33 | 2.1 | Institutional Delivery | 44.9 |
| 18 | Institutional Delivery against reported delivery ** | 1,543 | 98.8 | 1,521 | 97.9 | Home Delivery by SBA | 6.4 |
| 19 | Total reported Deliveries | 1,562 | | 1,554 | | PNC within 14 days of delivery | 28.8 |
| 20 | Reported Delivery at home assisted by a Doctor/Nurse /LHV/ANM(%) | 13 | 68.4 | 15 | 45.5 | | |
| 21 | Woman getting post partum check up between 48 hours and 14 days to total reported deliveries | 196 | 12.5 | 304 | 19.6 | | |
| Note: In case of HMIS data, Percentage is with reference to total number of Pregnant Women registered for ANC during the period and in DLHS-4 is with ref. to Women who had live/still births since three years preceding the survey. (DLHS-4 reference period is from 1-1-2008 to survey date). | | | | | | | |
| 22 | Expected number of Live Birth | 3,002 | | 3,036 | | | |
| D) Child Immunization : | | HMIS (2015-16) | | HMIS (2016-17) | | DLHS-4 (2012-13) | |
| 23 | Total reported live births | 1,570 | 52.3 | 1,659 | 54.7 | Breastfeed within 1 hour of birth | 49.3 |
| 24 | Total Still Birth | 16 | 0.5 | 57 | 1.9 | BCG | 67.4 |

| | | | | | | | |
|---|--|------------------------|------|------------------------|------|---------------------------|------|
| 25 | Newborn weighted at birth to live birth | 1,552 | 98.9 | 1,585 | 95.5 | 3 dose of DPT | 69.8 |
| 26 | Newborn having weight less than 2.5 kg to newborn weighted at birth. | 185 | 11.9 | 147 | 9.3 | Measles | 58.1 |
| 27 | Newborns breastfed within 1 hour to total reported live birth. | 1,519 | 96.8 | 1,494 | 90.1 | Full Immunization | 43 |
| 28 | Fully immunized | 1,733 | 57.7 | 1,857 | 61.2 | | |
| 29 | Children who have received BCG | 2,585 | 86.1 | 2,612 | 86.0 | | |
| 30 | Children who have received DPT1 | 1,761 | 58.7 | 218 | 7.2 | | |
| 31 | Children who have received DPT3 | 1,949 | 64.9 | 228 | 7.5 | | |
| 32 | Children who have received Measles | 2,021 | 67.3 | 2,016 | 66.4 | | |
| 33 | Diarrhoea and Dehydration cases reported in children below 5 years | 1,548 | | 1,622 | | | |
| <p>Note: In case of HMIS data, Proportions are calculated for 0-11 months old infants to total number of reported live births and in DLHS-4, the percentage of children 12-23 months olds who received specific vaccination to mothers who had last and last but one living child born three years preceding the survey. (DLHS-4 reference period is from 1-1-2008 to survey date).</p> | | | | | | | |
| E) Family Planning | | HMIS (2015-16) | | HMIS (2016-17) | | DLHS-4 (2012-13) | |
| 34 | Number of Vasectomies performed | 0 | | 0 | | Limiting | 18.5 |
| 35 | Number of Tubectomies performed | 67 | | 35 | | Spacing | 18.2 |
| 36 | Number of IUD insertion | 185 | | 179 | | Women using any method | 43.7 |
| 37 | Number of IUD removals | 99 | | 116 | | Women using modern method | 42 |
| F) Service delivery | | HMIS (2015-16) | | HMIS (2016-17) | | | |
| 38 | IPD cases per 10,000 population *** | 220 | | 255 | | | |
| 39 | General OPD cases per 10,000 population *** | 7756 | | 7457 | | | |
| 40 | Ayush OPD cases per 10,000 population *** | 588 | | 527 | | | |
| G) Mortality Details | | | | | | | |
| I) | Infant Deaths (In Numbers) | HMIS (2015-16) | | HMIS (2016-17) | | | |
| 41 | Neonatal death within 24 hours | 0 | | 1 | | | |
| 42 | Up to 1 Weeks of Birth | 0 | | 0 | | | |
| 43 | Between 1 week & 4 weeks of birth | 0 | | 0 | | | |
| 44 | Neonatal Death | 0 | | 1 | | | |
| 45 | Between 1 month and 11 months | 0 | | 0 | | | |
| 46 | Total reported infant deaths | 0 | | 1 | | | |
| II) | Maternal Mortality | | | | | | |
| | Maternal Deaths (In Numbers) | HMIS (2015-16) | | HMIS (2016-17) | | | |
| 47 | Due to Abortion | 0 | | 0 | | | |
| 48 | Due to Obstructed/prolonged labour | 0 | | 0 | | | |
| 49 | Due to Severe hypertension/fits | 0 | | 0 | | | |
| 50 | Due to Bleeding | 1 | | 0 | | | |
| 51 | Due to High fever | 0 | | 0 | | | |
| 52 | Due to Other Causes (including causes not known) | 0 | | 0 | | | |
| 53 | Total reported maternal deaths | 1 | | 0 | | | |

Longding Factsheet

Key Indicators

| A) Demographic Indicators (Census 2011) | | | | | | | |
|--|--|-----------------|----|-----------------|------|-----------------------------------|------|
| 1 | Population 2011 | 53,014 | | | | | |
| 2 | Sex Ratio | 931 | | | | | |
| B) Infrastructure Details (RHS 2015) | | Numbers | | | | | |
| 3 | Sub Centers (SC) | NA | | | | | |
| 4 | Primary Health Centers (PHC) | NA | | | | | |
| 5 | Community Health Centers (CHC) | NA | | | | | |
| 6 | SDH | NA | | | | | |
| 7 | DH | NA | | | | | |
| C) Maternal Health: | | 2015-16 | | 2016-17 | | | |
| 8 | Expected number of PW | 1,131 | | 1,133 | | | |
| 9 | Expected number of Delivery | 1,049 | | 1,050 | | | |
| | | HMIS (2015-16) | | HMIS (2016-17) | | DLHS-4 (2012-13) | |
| | | Number | PC | Number | PC | | |
| 10 | PW register for ANC | NA | NA | 736 | 65.0 | Any ANC | 43.4 |
| 11 | Mothers registered in the first trimester | NA | NA | 259 | 35.2 | ANC Check up in first trimester | 22.5 |
| 12 | Mothers who had at least 3 Ante-Natal care check ups | NA | NA | 266 | 36.1 | 3 or more ANC check up | 26.3 |
| 13 | PW who got TT 2 / Booster injection | NA | NA | 548 | 74.5 | At least 1 TT received | 43.1 |
| 14 | PW who consumed 100 IFA tablets | NA | NA | 567 | 77.0 | 100 IFA tablets consumed | 17.9 |
| 15 | Home Delivery against expected delivery * | NA | NA | 198 | 18.9 | Full ANC | 13.3 |
| 16 | Institutional Delivery against expected delivery * | NA | NA | 306 | 29.1 | Home Delivery | 68.8 |
| 17 | Home Delivery against reported delivery ** | NA | NA | 198 | 39.3 | Institutional Delivery | 29.9 |
| 18 | Institutional Delivery against reported delivery ** | NA | NA | 306 | 60.7 | Home Delivery by SBA | 6.4 |
| 19 | Total reported Deliveries | NA | | 504 | | PNC within 14 days of delivery | 42.0 |
| 20 | Reported Delivery at home assisted by a Doctor/Nurse /LHV/ANM(%) | NA | NA | 13 | 6.6 | | |
| 21 | Woman getting post partum check up between 48 hours and 14 days to total reported deliveries | NA | NA | 50 | 9.9 | | |
| Note: In case of HMIS data, Percentage is with reference to total number of Pregnant Women registered for ANC during the period and in DLHS-4 is with ref. to Women who had live/still births since three years preceding the survey. (DLHS-4 reference period is from 1-1-2008 to survey date). | | | | | | | |
| 22 | Expected number of Live Birth | 1,028 | | 1,030 | | | |
| D) Child Immunization : | | HMIS (2015-16) | | HMIS (2016-17) | | DLHS-4 (2012-13) | |
| 23 | Total reported live births | NA | NA | 503 | 48.8 | Breastfeed within 1 hour of birth | 47.7 |
| 24 | Total Still Birth | NA | NA | 2 | 0.2 | BCG | 60.2 |

| | | | | | | | |
|---|--|------------------------|----|------------------------|------|---------------------------|------|
| 25 | Newborn weighted at birth to live birth | NA | NA | 414 | 82.3 | 3 dose of DPT | 67 |
| 26 | Newborn having weight less than 2.5 kg to newborn weighted at birth. | NA | NA | 2 | 0.5 | Measles | 43.6 |
| 27 | Newborns breastfed within 1 hour to total reported live birth. | NA | NA | 420 | 83.5 | Full Immunization | 33.3 |
| 28 | Fully immunized | NA | NA | 461 | 44.8 | | |
| 29 | Children who have received BCG | NA | NA | 608 | 59.0 | | |
| 30 | Children who have received DPT1 | NA | NA | 3 | 0.3 | | |
| 31 | Children who have received DPT3 | NA | NA | 30 | 2.9 | | |
| 32 | Children who have received Measles | NA | NA | 480 | 46.6 | | |
| 33 | Diarrhoea and Dehydration cases reported in children below 5 years | NA | | 1,275 | | | |
| <p>Note: In case of HMIS data, Proportions are calculated for 0-11 months old infants to total number of reported live births and in DLHS-4, the percentage of children 12-23 months olds who received specific vaccination to mothers who had last and last but one living child born three years preceding the survey. (DLHS-4 reference period is from 1-1-2008 to survey date).</p> | | | | | | | |
| E) Family Planning | | HMIS (2015-16) | | HMIS (2016-17) | | DLHS-4 (2012-13) | |
| 34 | Number of Vasectomies performed | NA | | 0 | | Limiting | 11.6 |
| 35 | Number of Tubectomies performed | NA | | 0 | | Spacing | 22.6 |
| 36 | Number of IUD insertion | NA | | 3 | | Women using any method | 45.3 |
| 37 | Number of IUD removals | NA | | 9 | | Women using modern method | 42.8 |
| F) Service delivery | | HMIS (2015-16) | | HMIS (2016-17) | | | |
| 38 | IPD cases per 10,000 population *** | NA | | 146 | | | |
| 39 | General OPD cases per 10,000 population *** | NA | | 6629 | | | |
| 40 | Ayush OPD cases per 10,000 population *** | NA | | 2604 | | | |
| G) Mortality Details | | | | | | | |
| I) | Infant Deaths (In Numbers) | HMIS (2015-16) | | HMIS (2016-17) | | | |
| 41 | Neonatal death within 24 hours | NA | | 0 | | | |
| 42 | Up to 1 Weeks of Birth | NA | | 0 | | | |
| 43 | Between 1 week & 4 weeks of birth | NA | | 0 | | | |
| 44 | Neonatal Death | 0 | | 0 | | | |
| 45 | Between 1 month and 11 months | NA | | 0 | | | |
| 46 | Total reported infant deaths | 0 | | 0 | | | |
| II) | Maternal Mortality | HMIS (2015-16) | | HMIS (2016-17) | | | |
| | Maternal Deaths (In Numbers) | HMIS (2015-16) | | HMIS (2016-17) | | | |
| 47 | Due to Abortion | NA | | 0 | | | |
| 48 | Due to Obstructed/prolonged labour | NA | | 0 | | | |
| 49 | Due to Severe hypertension/fits | NA | | 0 | | | |
| 50 | Due to Bleeding | NA | | 0 | | | |
| 51 | Due to High fever | NA | | 0 | | | |
| 52 | Due to Other Causes (including causes not known) | NA | | 0 | | | |
| 53 | Total reported maternal deaths | 0 | | 0 | | | |

Lower Dibang Valley Factsheet

Key Indicators

| A) Demographic Indicators (Census 2011) | | | | | | | |
|--|--|----------------|-------|----------------|-------|-----------------------------------|------|
| 1 | Population 2011 | 53,986 | | | | | |
| 2 | Sex Ratio | 919 | | | | | |
| B) Infrastructure Details (RHS 2015) | | Numbers | | | | | |
| 3 | Sub Centers (SC) | NA | | | | | |
| 4 | Primary Health Centers (PHC) | NA | | | | | |
| 5 | Community Health Centers (CHC) | 3 | | | | | |
| 6 | SDH | 0 | | | | | |
| 7 | DH | 1 | | | | | |
| C) Maternal Health: | | 2015-16 | | 2016-17 | | | |
| 8 | Expected number of PW | 1,181 | | 1,183 | | | |
| 9 | Expected number of Delivery | 1,095 | | 1,097 | | | |
| | | HMIS (2015-16) | | HMIS (2016-17) | | DLHS-4 (2012-13) | |
| | | Number | PC | Number | PC | | |
| 10 | PW register for ANC | 1,183 | 100.2 | 1,282 | 108.4 | Any ANC | 72.4 |
| 11 | Mothers registered in the first trimester | 338 | 28.6 | 478 | 37.3 | ANC Check up in first trimester | 35.4 |
| 12 | Mothers who had at least 3 Ante-Natal care check ups | 595 | 50.3 | 752 | 58.7 | 3 or more ANC check up | 53.9 |
| 13 | PW who got TT 2 / Booster injection | 664 | 56.1 | 765 | 59.7 | At least 1 TT received | 70.0 |
| 14 | PW who consumed 100 IFA tablets | 583 | 49.3 | 833 | 65.0 | 100 IFA tablets consumed | 13.2 |
| 15 | Home Delivery against expected delivery * | 70 | 6.4 | 13 | 1.2 | Full ANC | 6.2 |
| 16 | Institutional Delivery against expected delivery * | 516 | 47.1 | 644 | 58.7 | Home Delivery | 38.5 |
| 17 | Home Delivery against reported delivery ** | 70 | 11.9 | 13 | 2.0 | Institutional Delivery | 60.3 |
| 18 | Institutional Delivery against reported delivery ** | 516 | 88.1 | 644 | 98.0 | Home Delivery by SBA | 2.6 |
| 19 | Total reported Deliveries | 586 | | 657 | | PNC within 14 days of delivery | 41.1 |
| 20 | Reported Delivery at home assisted by a Doctor/Nurse /LHV/ANM(%) | 59 | 84.3 | 13 | 100.0 | | |
| 21 | Woman getting post partum check up between 48 hours and 14 days to total reported deliveries | 351 | 59.9 | 163 | 24.8 | | |
| Note: In case of HMIS data, Percentage is with reference to total number of Pregnant Women registered for ANC during the period and in DLHS-4 is with ref. to Women who had live/still births since three years preceding the survey. (DLHS-4 reference period is from 1-1-2008 to survey date). | | | | | | | |
| 22 | Expected number of Live Birth | 1,073 | | 1,075 | | | |
| D) Child Immunization : | | HMIS (2015-16) | | HMIS (2016-17) | | DLHS-4 (2012-13) | |
| 23 | Total reported live births | 606 | 56.5 | 662 | 61.6 | Breastfeed within 1 hour of birth | 70.3 |
| 24 | Total Still Birth | 16 | 1.5 | 10 | 0.9 | BCG | 84.2 |

| | | | | | | | |
|---|--|------------------------|------|------------------------|------|---------------------------|------|
| 25 | Newborn weighted at birth to live birth | 601 | 99.2 | 657 | 99.2 | 3 dose of DPT | 73.7 |
| 26 | Newborn having weight less than 2.5 kg to newborn weighted at birth. | 61 | 10.1 | 57 | 8.7 | Measles | 73.7 |
| 27 | Newborns breastfed within 1 hour to total reported live birth. | 591 | 97.5 | 652 | 98.5 | Full Immunization | 64.9 |
| 28 | Fully immunized | 715 | 66.6 | 773 | 71.9 | | |
| 29 | Children who have received BCG | 849 | 79.1 | 918 | 85.4 | | |
| 30 | Children who have received DPT1 | 636 | 59.3 | 69 | 6.4 | | |
| 31 | Children who have received DPT3 | 654 | 60.9 | 92 | 8.6 | | |
| 32 | Children who have received Measles | 750 | 69.9 | 841 | 78.2 | | |
| 33 | Diarrhoea and Dehydration cases reported in children below 5 years | 651 | | 452 | | | |
| <p>Note: In case of HMIS data, Proportions are calculated for 0-11 months old infants to total number of reported live births and in DLHS-4, the percentage of children 12-23 months olds who received specific vaccination to mothers who had last and last but one living child born three years preceding the survey. (DLHS-4 reference period is from 1-1-2008 to survey date).</p> | | | | | | | |
| E) Family Planning | | HMIS (2015-16) | | HMIS (2016-17) | | DLHS-4 (2012-13) | |
| 34 | Number of Vasectomies performed | 2 | | 1 | | Limiting | 11.3 |
| 35 | Number of Tubectomies performed | 26 | | 21 | | Spacing | 20.1 |
| 36 | Number of IUD insertion | 55 | | 37 | | Women using any method | 51.7 |
| 37 | Number of IUD removals | 24 | | 15 | | Women using modern method | 40.1 |
| F) Service delivery | | HMIS (2015-16) | | HMIS (2016-17) | | | |
| 38 | IPD cases per 10,000 population *** | 398 | | 383 | | | |
| 39 | General OPD cases per 10,000 population *** | 9927 | | 11650 | | | |
| 40 | Ayush OPD cases per 10,000 population *** | 41 | | 41 | | | |
| G) Mortality Details | | | | | | | |
| I) | Infant Deaths (In Numbers) | HMIS (2015-16) | | HMIS (2016-17) | | | |
| 41 | Neonatal death within 24 hours | 2 | | 1 | | | |
| 42 | Up to 1 Weeks of Birth | 0 | | 2 | | | |
| 43 | Between 1 week & 4 weeks of birth | 0 | | 0 | | | |
| 44 | Neonatal Death | 2 | | 3 | | | |
| 45 | Between 1 month and 11 months | 5 | | 2 | | | |
| 46 | Total reported infant deaths | 7 | | 5 | | | |
| II) | Maternal Mortality | HMIS (2015-16) | | HMIS (2016-17) | | | |
| | Maternal Deaths (In Numbers) | HMIS (2015-16) | | HMIS (2016-17) | | | |
| 47 | Due to Abortion | 0 | | 0 | | | |
| 48 | Due to Obstructed/prolonged labour | 0 | | 0 | | | |
| 49 | Due to Severe hypertension/fits | 0 | | 0 | | | |
| 50 | Due to Bleeding | 1 | | 0 | | | |
| 51 | Due to High fever | 0 | | 0 | | | |
| 52 | Due to Other Causes (including causes not known) | 0 | | 0 | | | |
| 53 | Total reported maternal deaths | 1 | | 0 | | | |

Lower Subansiri Factsheet

Key Indicators

| A) Demographic Indicators (Census 2011) | | | | | | | |
|--|--|----------------|------|----------------|------|-----------------------------------|------|
| 1 | Population 2011 | 82,839 | | | | | |
| 2 | Sex Ratio | 975 | | | | | |
| B) Infrastructure Details (RHS 2015) | | Numbers | | | | | |
| 3 | Sub Centers (SC) | NA | | | | | |
| 4 | Primary Health Centers (PHC) | NA | | | | | |
| 5 | Community Health Centers (CHC) | 2 | | | | | |
| 6 | SDH | 0 | | | | | |
| 7 | DH | 1 | | | | | |
| C) Maternal Health: | | 2015-16 | | 2016-17 | | | |
| 8 | Expected number of PW | 2,132 | | 2,224 | | | |
| 9 | Expected number of Delivery | 1,977 | | 2,062 | | | |
| | | HMIS (2015-16) | | HMIS (2016-17) | | DLHS-4 (2012-13) | |
| | | Number | PC | Number | PC | | |
| 10 | PW register for ANC | 1,225 | 57.5 | 1,611 | 72.4 | Any ANC | 79.3 |
| 11 | Mothers registered in the first trimester | 377 | 30.8 | 428 | 26.6 | ANC Check up in first trimester | 48.3 |
| 12 | Mothers who had at least 3 Ante-Natal care check ups | 402 | 32.8 | 435 | 27.0 | 3 or more ANC check up | 60.1 |
| 13 | PW who got TT 2 / Booster injection | 570 | 46.5 | 535 | 33.2 | At least 1 TT received | 78.6 |
| 14 | PW who consumed 100 IFA tablets | 941 | 76.8 | 835 | 51.8 | 100 IFA tablets consumed | 25.1 |
| 15 | Home Delivery against expected delivery * | 121 | 6.1 | 154 | 7.5 | Full ANC | 22.3 |
| 16 | Institutional Delivery against expected delivery * | 485 | 24.5 | 427 | 20.7 | Home Delivery | 33.2 |
| 17 | Home Delivery against reported delivery ** | 121 | 20.0 | 154 | 26.5 | Institutional Delivery | 65.9 |
| 18 | Institutional Delivery against reported delivery ** | 485 | 80.0 | 427 | 73.5 | Home Delivery by SBA | 4.3 |
| 19 | Total reported Deliveries | 606 | | 581 | | PNC within 14 days of delivery | 46.0 |
| 20 | Reported Delivery at home assisted by a Doctor/Nurse /LHV/ANM(%) | 22 | 18.2 | 17 | 11.0 | | |
| 21 | Woman getting post partum check up between 48 hours and 14 days to total reported deliveries | 187 | 30.9 | 189 | 32.5 | | |
| Note: In case of HMIS data, Percentage is with reference to total number of Pregnant Women registered for ANC during the period and in DLHS-4 is with ref. to Women who had live/still births since three years preceding the survey. (DLHS-4 reference period is from 1-1-2008 to survey date). | | | | | | | |
| 22 | Expected number of Live Birth | 1,938 | | 2,022 | | | |
| D) Child Immunization : | | HMIS (2015-16) | | HMIS (2016-17) | | DLHS-4 (2012-13) | |
| 23 | Total reported live births | 486 | 25.1 | 582 | 28.8 | Breastfeed within 1 hour of birth | 69.2 |
| 24 | Total Still Birth | 11 | 0.6 | 13 | 0.6 | BCG | 89 |

| | | | | | | | |
|---|--|------------------------|-------|------------------------|------|---------------------------|------|
| 25 | Newborn weighted at birth to live birth | 487 | 100.2 | 474 | 81.4 | 3 dose of DPT | 83.6 |
| 26 | Newborn having weight less than 2.5 kg to newborn weighted at birth. | 22 | 4.5 | 50 | 10.5 | Measles | 76.7 |
| 27 | Newborns breastfed within 1 hour to total reported live birth. | 429 | 88.3 | 413 | 71.0 | Full Immunization | 68.5 |
| 28 | Fully immunized | 751 | 38.8 | 813 | 40.2 | | |
| 29 | Children who have received BCG | 713 | 36.8 | 610 | 30.2 | | |
| 30 | Children who have received DPT1 | 689 | 35.6 | 54 | 2.7 | | |
| 31 | Children who have received DPT3 | 743 | 38.3 | 64 | 3.2 | | |
| 32 | Children who have received Measles | 828 | 42.7 | 895 | 44.3 | | |
| 33 | Diarrhoea and Dehydration cases reported in children below 5 years | 1,217 | | 1,475 | | | |
| <p>Note: In case of HMIS data, Proportions are calculated for 0-11 months old infants to total number of reported live births and in DLHS-4, the percentage of children 12-23 months olds who received specific vaccination to mothers who had last and last but one living child born three years preceding the survey. (DLHS-4 reference period is from 1-1-2008 to survey date).</p> | | | | | | | |
| E) Family Planning | | HMIS (2015-16) | | HMIS (2016-17) | | DLHS-4 (2012-13) | |
| 34 | Number of Vasectomies performed | 0 | | 0 | | Limiting | 7.4 |
| 35 | Number of Tubectomies performed | 2 | | 7 | | Spacing | 11.5 |
| 36 | Number of IUD insertion | 205 | | 160 | | Women using any method | 57.4 |
| 37 | Number of IUD removals | 131 | | 68 | | Women using modern method | 50.6 |
| F) Service delivery | | HMIS (2015-16) | | HMIS (2016-17) | | | |
| 38 | IPD cases per 10,000 population *** | 359 | | 357 | | | |
| 39 | General OPD cases per 10,000 population *** | 5175 | | 4913 | | | |
| 40 | Ayush OPD cases per 10,000 population *** | 208 | | 403 | | | |
| G) Mortality Details | | | | | | | |
| I) | Infant Deaths (In Numbers) | HMIS (2015-16) | | HMIS (2016-17) | | | |
| 41 | Neonatal death within 24 hours | 0 | | 0 | | | |
| 42 | Up to 1 Weeks of Birth | 2 | | 0 | | | |
| 43 | Between 1 week & 4 weeks of birth | 0 | | 0 | | | |
| 44 | Neonatal Death | 2 | | 0 | | | |
| 45 | Between 1 month and 11 months | 1 | | 3 | | | |
| 46 | Total reported infant deaths | 3 | | 3 | | | |
| II) | Maternal Mortality | HMIS (2015-16) | | HMIS (2016-17) | | | |
| | Maternal Deaths (In Numbers) | HMIS (2015-16) | | HMIS (2016-17) | | | |
| 47 | Due to Abortion | 0 | | 0 | | | |
| 48 | Due to Obstructed/prolonged labour | 0 | | 0 | | | |
| 49 | Due to Severe hypertension/fits | 0 | | 0 | | | |
| 50 | Due to Bleeding | 0 | | 0 | | | |
| 51 | Due to High fever | 1 | | 0 | | | |
| 52 | Due to Other Causes (including causes not known) | 0 | | 0 | | | |
| 53 | Total reported maternal deaths | 1 | | 0 | | | |

Papum Pare Factsheet

Key Indicators

| A) Demographic Indicators (Census 2011) | | | | | | | |
|--|--|----------------|-------|----------------|-------|-----------------------------------|------|
| 1 | Population 2011 | 176,385 | | | | | |
| 2 | Sex Ratio | 950 | | | | | |
| B) Infrastructure Details (RHS 2015) | | Numbers | | | | | |
| 3 | Sub Centers (SC) | NA | | | | | |
| 4 | Primary Health Centers (PHC) | NA | | | | | |
| 5 | Community Health Centers (CHC) | 3 | | | | | |
| 6 | SDH | 0 | | | | | |
| 7 | DH | 1 | | | | | |
| C) Maternal Health: | | 2015-16 | | 2016-17 | | | |
| 8 | Expected number of PW | 4,463 | | 4,638 | | | |
| 9 | Expected number of Delivery | 4,138 | | 4,301 | | | |
| | | HMIS (2015-16) | | HMIS (2016-17) | | DLHS-4 (2012-13) | |
| | | Number | PC | Number | PC | | |
| 10 | PW register for ANC | 4,263 | 95.5 | 6,045 | 130.3 | Any ANC | 78.0 |
| 11 | Mothers registered in the first trimester | 2,035 | 47.7 | 2,008 | 33.2 | ANC Check up in first trimester | 47.8 |
| 12 | Mothers who had at least 3 Ante-Natal care check ups | 2,255 | 52.9 | 1,756 | 29.0 | 3 or more ANC check up | 55.2 |
| 13 | PW who got TT 2 / Booster injection | 992 | 23.3 | 1,018 | 16.8 | At least 1 TT received | 76.0 |
| 14 | PW who consumed 100 IFA tablets | 1,032 | 24.2 | 2,416 | 40.0 | 100 IFA tablets consumed | 29.4 |
| 15 | Home Delivery against expected delivery * | 30 | 0.7 | 20 | 0.5 | Full ANC | 22.2 |
| 16 | Institutional Delivery against expected delivery * | 4,407 | 106.5 | 4,416 | 102.7 | Home Delivery | 22.7 |
| 17 | Home Delivery against reported delivery ** | 30 | 0.7 | 20 | 0.5 | Institutional Delivery | 74.1 |
| 18 | Institutional Delivery against reported delivery ** | 4,407 | 99.3 | 4,416 | 99.5 | Home Delivery by SBA | 0.9 |
| 19 | Total reported Deliveries | 4,437 | | 4,436 | | PNC within 14 days of delivery | 52.1 |
| 20 | Reported Delivery at home assisted by a Doctor/Nurse /LHV/ANM(%) | 29 | 96.7 | 8 | 40.0 | | |
| 21 | Woman getting post partum check up between 48 hours and 14 days to total reported deliveries | 483 | 10.9 | 326 | 7.3 | | |
| Note: In case of HMIS data, Percentage is with reference to total number of Pregnant Women registered for ANC during the period and in DLHS-4 is with ref. to Women who had live/still births since three years preceding the survey. (DLHS-4 reference period is from 1-1-2008 to survey date). | | | | | | | |
| 22 | Expected number of Live Birth | 4,057 | | 4,216 | | | |
| D) Child Immunization : | | HMIS (2015-16) | | HMIS (2016-17) | | DLHS-4 (2012-13) | |
| 23 | Total reported live births | 4,598 | 113.3 | 4,717 | 111.9 | Breastfeed within 1 hour of birth | 44.9 |
| 24 | Total Still Birth | 77 | 1.9 | 51 | 1.2 | BCG | 93 |

| | | | | | | | |
|---|--|------------------------|-------|------------------------|-------|---------------------------|------|
| 25 | Newborn weighted at birth to live birth | 3,794 | 82.5 | 4,731 | 100.3 | 3 dose of DPT | 72.7 |
| 26 | Newborn having weight less than 2.5 kg to newborn weighted at birth. | 214 | 5.6 | 537 | 11.4 | Measles | 74.7 |
| 27 | Newborns breastfed within 1 hour to total reported live birth. | 4,423 | 96.2 | 3,757 | 79.6 | Full Immunization | 65.5 |
| 28 | Fully immunized | 3,099 | 76.4 | 4,281 | 101.5 | | |
| 29 | Children who have received BCG | 4,198 | 103.5 | 4,029 | 95.6 | | |
| 30 | Children who have received DPT1 | 3,101 | 76.4 | 352 | 8.3 | | |
| 31 | Children who have received DPT3 | 2,853 | 70.3 | 269 | 6.4 | | |
| 32 | Children who have received Measles | 3,252 | 80.2 | 2,993 | 71.0 | | |
| 33 | Diarrhoea and Dehydration cases reported in children below 5 years | 2,781 | | 1,721 | | | |
| <p>Note: In case of HMIS data, Proportions are calculated for 0-11 months old infants to total number of reported live births and in DLHS-4, the percentage of children 12-23 months olds who received specific vaccination to mothers who had last and last but one living child born three years preceding the survey. (DLHS-4 reference period is from 1-1-2008 to survey date).</p> | | | | | | | |
| E) Family Planning | | HMIS (2015-16) | | HMIS (2016-17) | | DLHS-4 (2012-13) | |
| 34 | Number of Vasectomies performed | 0 | | 28 | | Limiting | 10.8 |
| 35 | Number of Tubectomies performed | 320 | | 412 | | Spacing | 21.3 |
| 36 | Number of IUD insertion | 526 | | 577 | | Women using any method | 49.3 |
| 37 | Number of IUD removals | 92 | | 149 | | Women using modern method | 43.3 |
| F) Service delivery | | HMIS (2015-16) | | HMIS (2016-17) | | | |
| 38 | IPD cases per 10,000 population *** | 1389 | | 1052 | | | |
| 39 | General OPD cases per 10,000 population *** | 15803 | | 15291 | | | |
| 40 | Ayush OPD cases per 10,000 population *** | 264 | | 190 | | | |
| G) Mortality Details | | | | | | | |
| I) | Infant Deaths (In Numbers) | HMIS (2015-16) | | HMIS (2016-17) | | | |
| 41 | Neonatal death within 24 hours | 2 | | 6 | | | |
| 42 | Up to 1 Weeks of Birth | 17 | | 17 | | | |
| 43 | Between 1 week & 4 weeks of birth | 0 | | 0 | | | |
| 44 | Neonatal Death | 19 | | 23 | | | |
| 45 | Between 1 month and 11 months | 11 | | 6 | | | |
| 46 | Total reported infant deaths | 30 | | 29 | | | |
| II) | Maternal Mortality | HMIS (2015-16) | | HMIS (2016-17) | | | |
| | Maternal Deaths (In Numbers) | HMIS (2015-16) | | HMIS (2016-17) | | | |
| 47 | Due to Abortion | 0 | | 0 | | | |
| 48 | Due to Obstructed/prolonged labour | 2 | | 0 | | | |
| 49 | Due to Severe hypertension/fits | 0 | | 0 | | | |
| 50 | Due to Bleeding | 0 | | 0 | | | |
| 51 | Due to High fever | 0 | | 0 | | | |
| 52 | Due to Other Causes (including causes not known) | 2 | | 1 | | | |
| 53 | Total reported maternal deaths | 4 | | 1 | | | |

Tawang Factsheet

Key Indicators

| A) Demographic Indicators (Census 2011) | | | | | | | |
|--|--|----------------|------|----------------|------|-----------------------------------|------|
| 1 | Population 2011 | 49,950 | | | | | |
| 2 | Sex Ratio | 701 | | | | | |
| B) Infrastructure Details (RHS 2015) | | Numbers | | | | | |
| 3 | Sub Centers (SC) | NA | | | | | |
| 4 | Primary Health Centers (PHC) | NA | | | | | |
| 5 | Community Health Centers (CHC) | 1 | | | | | |
| 6 | SDH | 0 | | | | | |
| 7 | DH | 1 | | | | | |
| C) Maternal Health: | | 2015-16 | | 2016-17 | | | |
| 8 | Expected number of PW | 1,186 | | 1,214 | | | |
| 9 | Expected number of Delivery | 1,100 | | 1,125 | | | |
| | | HMIS (2015-16) | | HMIS (2016-17) | | DLHS-4 (2012-13) | |
| | | Number | PC | Number | PC | | |
| 10 | PW register for ANC | 881 | 74.3 | 806 | 66.4 | Any ANC | 61.5 |
| 11 | Mothers registered in the first trimester | 404 | 45.9 | 289 | 35.9 | ANC Check up in first trimester | 36.7 |
| 12 | Mothers who had at least 3 Ante-Natal care check ups | 225 | 25.5 | 232 | 28.8 | 3 or more ANC check up | 42.7 |
| 13 | PW who got TT 2 / Booster injection | 438 | 49.7 | 497 | 61.7 | At least 1 TT received | 58.8 |
| 14 | PW who consumed 100 IFA tablets | 505 | 57.3 | 570 | 70.7 | 100 IFA tablets consumed | 17.4 |
| 15 | Home Delivery against expected delivery * | 84 | 7.6 | 113 | 10.0 | Full ANC | 13.5 |
| 16 | Institutional Delivery against expected delivery * | 342 | 31.1 | 337 | 29.9 | Home Delivery | 62.3 |
| 17 | Home Delivery against reported delivery ** | 84 | 19.7 | 113 | 25.1 | Institutional Delivery | 37.0 |
| 18 | Institutional Delivery against reported delivery ** | 342 | 80.3 | 337 | 74.9 | Home Delivery by SBA | 4.2 |
| 19 | Total reported Deliveries | 426 | | 450 | | PNC within 14 days of delivery | 53.1 |
| 20 | Reported Delivery at home assisted by a Doctor/Nurse /LHV/ANM(%) | 53 | 63.1 | 92 | 81.4 | | |
| 21 | Woman getting post partum check up between 48 hours and 14 days to total reported deliveries | 223 | 52.3 | 197 | 43.8 | | |
| Note: In case of HMIS data, Percentage is with reference to total number of Pregnant Women registered for ANC during the period and in DLHS-4 is with ref. to Women who had live/still births since three years preceding the survey. (DLHS-4 reference period is from 1-1-2008 to survey date). | | | | | | | |
| 22 | Expected number of Live Birth | 1,079 | | 1,103 | | | |
| D) Child Immunization : | | HMIS (2015-16) | | HMIS (2016-17) | | DLHS-4 (2012-13) | |
| 23 | Total reported live births | 426 | 39.5 | 444 | 40.2 | Breastfeed within 1 hour of birth | 62 |
| 24 | Total Still Birth | 7 | 0.6 | 8 | 0.7 | BCG | 83.3 |

| | | | | | | | |
|---|--|------------------------|------|------------------------|------|---------------------------|------|
| 25 | Newborn weighted at birth to live birth | 377 | 88.5 | 340 | 76.6 | 3 dose of DPT | 76.4 |
| 26 | Newborn having weight less than 2.5 kg to newborn weighted at birth. | 25 | 6.6 | 14 | 4.1 | Measles | 63.9 |
| 27 | Newborns breastfed within 1 hour to total reported live birth. | 361 | 84.7 | 390 | 87.8 | Full Immunization | 56.9 |
| 28 | Fully immunized | 635 | 58.9 | 639 | 57.9 | | |
| 29 | Children who have received BCG | 683 | 63.3 | 665 | 60.3 | | |
| 30 | Children who have received DPT1 | 489 | 45.3 | 0 | 0.0 | | |
| 31 | Children who have received DPT3 | 625 | 57.9 | 10 | 0.9 | | |
| 32 | Children who have received Measles | 651 | 60.4 | 671 | 60.8 | | |
| 33 | Diarrhoea and Dehydration cases reported in children below 5 years | 1,086 | | 1,600 | | | |
| <p>Note: In case of HMIS data, Proportions are calculated for 0-11 months old infants to total number of reported live births and in DLHS-4, the percentage of children 12-23 months olds who received specific vaccination to mothers who had last and last but one living child born three years preceding the survey. (DLHS-4 reference period is from 1-1-2008 to survey date).</p> | | | | | | | |
| E) Family Planning | | HMIS (2015-16) | | HMIS (2016-17) | | DLHS-4 (2012-13) | |
| 34 | Number of Vasectomies performed | 0 | | 0 | | Limiting | 14.1 |
| 35 | Number of Tubectomies performed | 0 | | 0 | | Spacing | 17.6 |
| 36 | Number of IUD insertion | 98 | | 80 | | Women using any method | 47.1 |
| 37 | Number of IUD removals | 57 | | 29 | | Women using modern method | 43.9 |
| F) Service delivery | | HMIS (2015-16) | | HMIS (2016-17) | | | |
| 38 | IPD cases per 10,000 population *** | 404 | | 404 | | | |
| 39 | General OPD cases per 10,000 population *** | 12001 | | 10930 | | | |
| 40 | Ayush OPD cases per 10,000 population *** | 545 | | 522 | | | |
| G) Mortality Details | | | | | | | |
| I) | Infant Deaths (In Numbers) | HMIS (2015-16) | | HMIS (2016-17) | | | |
| 41 | Neonatal death within 24 hours | 0 | | 0 | | | |
| 42 | Up to 1 Weeks of Birth | 1 | | 1 | | | |
| 43 | Between 1 week & 4 weeks of birth | 0 | | 0 | | | |
| 44 | Neonatal Death | 1 | | 1 | | | |
| 45 | Between 1 month and 11 months | 2 | | 1 | | | |
| 46 | Total reported infant deaths | 3 | | 2 | | | |
| II) | Maternal Mortality | HMIS (2015-16) | | HMIS (2016-17) | | | |
| | Maternal Deaths (In Numbers) | HMIS (2015-16) | | HMIS (2016-17) | | | |
| 47 | Due to Abortion | 0 | | 0 | | | |
| 48 | Due to Obstructed/prolonged labour | 0 | | 0 | | | |
| 49 | Due to Severe hypertension/fits | 0 | | 1 | | | |
| 50 | Due to Bleeding | 0 | | 0 | | | |
| 51 | Due to High fever | 0 | | 0 | | | |
| 52 | Due to Other Causes (including causes not known) | 0 | | 0 | | | |
| 53 | Total reported maternal deaths | 0 | | 1 | | | |

Tirap Factsheet

Key Indicators

| A) Demographic Indicators (Census 2011) | | | | | | | |
|--|--|-----------------|------|-----------------|------|-----------------------------------|------|
| 1 | Population 2011 | 58,961 | | | | | |
| 2 | Sex Ratio | 931 | | | | | |
| B) Infrastructure Details (RHS 2015) | | Numbers | | | | | |
| 3 | Sub Centers (SC) | NA | | | | | |
| 4 | Primary Health Centers (PHC) | NA | | | | | |
| 5 | Community Health Centers (CHC) | 1 | | | | | |
| 6 | SDH | 0 | | | | | |
| 7 | DH | 1 | | | | | |
| C) Maternal Health: | | 2015-16 | | 2016-17 | | | |
| 8 | Expected number of PW | 1,252 | | 1,260 | | | |
| 9 | Expected number of Delivery | 1,161 | | 1,168 | | | |
| | | HMIS (2015-16) | | HMIS (2016-17) | | DLHS-4 (2012-13) | |
| | | Number | PC | Number | PC | | |
| 10 | PW register for ANC | 800 | 63.9 | 1,055 | 83.7 | Any ANC | 43.4 |
| 11 | Mothers registered in the first trimester | 265 | 33.1 | 368 | 34.9 | ANC Check up in first trimester | 22.5 |
| 12 | Mothers who had at least 3 Ante-Natal care check ups | 350 | 43.8 | 410 | 38.9 | 3 or more ANC check up | 26.3 |
| 13 | PW who got TT 2 / Booster injection | 696 | 87.0 | 609 | 57.7 | At least 1 TT received | 43.1 |
| 14 | PW who consumed 100 IFA tablets | 499 | 62.4 | 239 | 22.7 | 100 IFA tablets consumed | 17.9 |
| 15 | Home Delivery against expected delivery * | 138 | 11.9 | 88 | 7.5 | Full ANC | 13.3 |
| 16 | Institutional Delivery against expected delivery * | 415 | 35.7 | 471 | 40.3 | Home Delivery | 68.8 |
| 17 | Home Delivery against reported delivery ** | 138 | 25.0 | 88 | 15.7 | Institutional Delivery | 29.9 |
| 18 | Institutional Delivery against reported delivery ** | 415 | 75.0 | 471 | 84.3 | Home Delivery by SBA | 6.4 |
| 19 | Total reported Deliveries | 553 | | 559 | | PNC within 14 days of delivery | 42.0 |
| 20 | Reported Delivery at home assisted by a Doctor/Nurse /LHV/ANM(%) | 10 | 7.2 | 6 | 6.8 | | |
| 21 | Woman getting post partum check up between 48 hours and 14 days to total reported deliveries | 232 | 42.0 | 340 | 60.8 | | |
| Note: In case of HMIS data, Percentage is with reference to total number of Pregnant Women registered for ANC during the period and in DLHS-4 is with ref. to Women who had live/still births since three years preceding the survey. (DLHS-4 reference period is from 1-1-2008 to survey date). | | | | | | | |
| 22 | Expected number of Live Birth | 1,138 | | 1,145 | | | |
| D) Child Immunization : | | HMIS (2015-16) | | HMIS (2016-17) | | DLHS-4 (2012-13) | |
| 23 | Total reported live births | 544 | 47.8 | 561 | 49.0 | Breastfeed within 1 hour of birth | 47.7 |
| 24 | Total Still Birth | 9 | 0.8 | 2 | 0.2 | BCG | 60.2 |

| | | | | | | | |
|---|--|------------------------|------|------------------------|-------|---------------------------|------|
| 25 | Newborn weighted at birth to live birth | 520 | 95.6 | 561 | 100.0 | 3 dose of DPT | 67 |
| 26 | Newborn having weight less than 2.5 kg to newborn weighted at birth. | 1 | 0.2 | 9 | 1.6 | Measles | 43.6 |
| 27 | Newborns breastfed within 1 hour to total reported live birth. | 520 | 95.6 | 560 | 99.8 | Full Immunization | 33.3 |
| 28 | Fully immunized | 580 | 51.0 | 563 | 49.2 | | |
| 29 | Children who have received BCG | 821 | 72.1 | 774 | 67.6 | | |
| 30 | Children who have received DPT1 | 636 | 55.9 | 3 | 0.3 | | |
| 31 | Children who have received DPT3 | 669 | 58.8 | 7 | 0.6 | | |
| 32 | Children who have received Measles | 616 | 54.1 | 668 | 58.3 | | |
| 33 | Diarrhoea and Dehydration cases reported in children below 5 years | 422 | | 399 | | | |
| <p>Note: In case of HMIS data, Proportions are calculated for 0-11 months old infants to total number of reported live births and in DLHS-4, the percentage of children 12-23 months olds who received specific vaccination to mothers who had last and last but one living child born three years preceding the survey. (DLHS-4 reference period is from 1-1-2008 to survey date).</p> | | | | | | | |
| E) Family Planning | | HMIS (2015-16) | | HMIS (2016-17) | | DLHS-4 (2012-13) | |
| 34 | Number of Vasectomies performed | 0 | | 0 | | Limiting | 11.6 |
| 35 | Number of Tubectomies performed | 29 | | 183 | | Spacing | 22.6 |
| 36 | Number of IUD insertion | 45 | | 44 | | Women using any method | 45.3 |
| 37 | Number of IUD removals | 47 | | 53 | | Women using modern method | 42.8 |
| F) Service delivery | | HMIS (2015-16) | | HMIS (2016-17) | | | |
| 38 | IPD cases per 10,000 population *** | 346 | | 411 | | | |
| 39 | General OPD cases per 10,000 population *** | 7255 | | 7003 | | | |
| 40 | Ayush OPD cases per 10,000 population *** | 1571 | | 1565 | | | |
| G) Mortality Details | | | | | | | |
| I) | Infant Deaths (In Numbers) | HMIS (2015-16) | | HMIS (2016-17) | | | |
| 41 | Neonatal death within 24 hours | 0 | | 2 | | | |
| 42 | Up to 1 Weeks of Birth | 0 | | 0 | | | |
| 43 | Between 1 week & 4 weeks of birth | 0 | | 0 | | | |
| 44 | Neonatal Death | 0 | | 2 | | | |
| 45 | Between 1 month and 11 months | 0 | | 0 | | | |
| 46 | Total reported infant deaths | 0 | | 2 | | | |
| II) | Maternal Mortality | HMIS (2015-16) | | HMIS (2016-17) | | | |
| | Maternal Deaths (In Numbers) | HMIS (2015-16) | | HMIS (2016-17) | | | |
| 47 | Due to Abortion | 0 | | 0 | | | |
| 48 | Due to Obstructed/prolonged labour | 0 | | 0 | | | |
| 49 | Due to Severe hypertension/fits | 1 | | 0 | | | |
| 50 | Due to Bleeding | 1 | | 0 | | | |
| 51 | Due to High fever | 0 | | 0 | | | |
| 52 | Due to Other Causes (including causes not known) | 0 | | 1 | | | |
| 53 | Total reported maternal deaths | 2 | | 1 | | | |

Upper Siang Factsheet

Key Indicators

| A) Demographic Indicators (Census 2011) | | | | | | | |
|--|--|-----------------|------|-----------------|------|-----------------------------------|------|
| 1 | Population 2011 | 35,289 | | | | | |
| 2 | Sex Ratio | 891 | | | | | |
| B) Infrastructure Details (RHS 2015) | | Numbers | | | | | |
| 3 | Sub Centers (SC) | NA | | | | | |
| 4 | Primary Health Centers (PHC) | NA | | | | | |
| 5 | Community Health Centers (CHC) | 4 | | | | | |
| 6 | SDH | 0 | | | | | |
| 7 | DH | 1 | | | | | |
| C) Maternal Health: | | 2015-16 | | 2016-17 | | | |
| 8 | Expected number of PW | 767 | | 768 | | | |
| 9 | Expected number of Delivery | 712 | | 712 | | | |
| | | HMIS (2015-16) | | HMIS (2016-17) | | DLHS-4 (2012-13) | |
| | | Number | PC | Number | PC | | |
| 10 | PW register for ANC | 733 | 95.5 | 603 | 78.5 | Any ANC | 28.4 |
| 11 | Mothers registered in the first trimester | 326 | 44.5 | 275 | 45.6 | ANC Check up in first trimester | 14.0 |
| 12 | Mothers who had at least 3 Ante-Natal care check ups | 383 | 52.3 | 343 | 56.9 | 3 or more ANC check up | 16.6 |
| 13 | PW who got TT 2 / Booster injection | 442 | 60.3 | 374 | 62.0 | At least 1 TT received | 25.4 |
| 14 | PW who consumed 100 IFA tablets | 478 | 65.2 | 581 | 96.4 | 100 IFA tablets consumed | 8.7 |
| 15 | Home Delivery against expected delivery * | 24 | 3.4 | 16 | 2.2 | Full ANC | 5.7 |
| 16 | Institutional Delivery against expected delivery * | 341 | 47.9 | 311 | 43.7 | Home Delivery | 66.4 |
| 17 | Home Delivery against reported delivery ** | 24 | 6.6 | 16 | 4.9 | Institutional Delivery | 30.1 |
| 18 | Institutional Delivery against reported delivery ** | 341 | 93.4 | 311 | 95.1 | Home Delivery by SBA | 4.4 |
| 19 | Total reported Deliveries | 365 | | 327 | | PNC within 14 days of delivery | 47.8 |
| 20 | Reported Delivery at home assisted by a Doctor/Nurse /LHV/ANM(%) | 17 | 70.8 | 13 | 81.3 | | |
| 21 | Woman getting post partum check up between 48 hours and 14 days to total reported deliveries | 106 | 29.0 | 181 | 55.4 | | |
| Note: In case of HMIS data, Percentage is with reference to total number of Pregnant Women registered for ANC during the period and in DLHS-4 is with ref. to Women who had live/still births since three years preceding the survey. (DLHS-4 reference period is from 1-1-2008 to survey date). | | | | | | | |
| 22 | Expected number of Live Birth | 698 | | 698 | | | |
| D) Child Immunization : | | HMIS (2015-16) | | HMIS (2016-17) | | DLHS-4 (2012-13) | |
| 23 | Total reported live births | 386 | 55.3 | 332 | 47.6 | Breastfeed within 1 hour of birth | 70.4 |
| 24 | Total Still Birth | 16 | 2.3 | 2 | 0.3 | BCG | 60 |

| | | | | | | | |
|---|--|------------------------|------|------------------------|------|---------------------------|------|
| 25 | Newborn weighted at birth to live birth | 380 | 98.4 | 298 | 89.8 | 3 dose of DPT | 50.9 |
| 26 | Newborn having weight less than 2.5 kg to newborn weighted at birth. | 23 | 6.1 | 22 | 7.4 | Measles | 40 |
| 27 | Newborns breastfed within 1 hour to total reported live birth. | 370 | 95.9 | 282 | 84.9 | Full Immunization | 29.1 |
| 28 | Fully immunized | 488 | 70.0 | 460 | 65.9 | | |
| 29 | Children who have received BCG | 507 | 72.7 | 311 | 44.6 | | |
| 30 | Children who have received DPT1 | 426 | 61.1 | 31 | 4.4 | | |
| 31 | Children who have received DPT3 | 492 | 70.5 | 33 | 4.7 | | |
| 32 | Children who have received Measles | 545 | 78.1 | 451 | 64.6 | | |
| 33 | Diarrhoea and Dehydration cases reported in children below 5 years | 604 | | 555 | | | |
| <p>Note: In case of HMIS data, Proportions are calculated for 0-11 months old infants to total number of reported live births and in DLHS-4, the percentage of children 12-23 months olds who received specific vaccination to mothers who had last and last but one living child born three years preceding the survey. (DLHS-4 reference period is from 1-1-2008 to survey date).</p> | | | | | | | |
| E) Family Planning | | HMIS (2015-16) | | HMIS (2016-17) | | DLHS-4 (2012-13) | |
| 34 | Number of Vasectomies performed | 0 | | 0 | | Limiting | 14.7 |
| 35 | Number of Tubectomies performed | 52 | | 25 | | Spacing | 28.9 |
| 36 | Number of IUD insertion | 117 | | 104 | | Women using any method | 41.7 |
| 37 | Number of IUD removals | 49 | | 36 | | Women using modern method | 39.8 |
| F) Service delivery | | HMIS (2015-16) | | HMIS (2016-17) | | | |
| 38 | IPD cases per 10,000 population *** | 403 | | 356 | | | |
| 39 | General OPD cases per 10,000 population *** | 12002 | | 11433 | | | |
| 40 | Ayush OPD cases per 10,000 population *** | 973 | | 902 | | | |
| G) Mortality Details | | | | | | | |
| I) | Infant Deaths (In Numbers) | HMIS (2015-16) | | HMIS (2016-17) | | | |
| 41 | Neonatal death within 24 hours | 0 | | 0 | | | |
| 42 | Up to 1 Weeks of Birth | 0 | | 0 | | | |
| 43 | Between 1 week & 4 weeks of birth | 0 | | 0 | | | |
| 44 | Neonatal Death | 0 | | 0 | | | |
| 45 | Between 1 month and 11 months | 0 | | 0 | | | |
| 46 | Total reported infant deaths | 0 | | 0 | | | |
| II) | Maternal Mortality | HMIS (2015-16) | | HMIS (2016-17) | | | |
| | Maternal Deaths (In Numbers) | HMIS (2015-16) | | HMIS (2016-17) | | | |
| 47 | Due to Abortion | 0 | | 0 | | | |
| 48 | Due to Obstructed/prolonged labour | 0 | | 0 | | | |
| 49 | Due to Severe hypertension/fits | 0 | | 0 | | | |
| 50 | Due to Bleeding | 0 | | 0 | | | |
| 51 | Due to High fever | 0 | | 0 | | | |
| 52 | Due to Other Causes (including causes not known) | 0 | | 0 | | | |
| 53 | Total reported maternal deaths | 0 | | 0 | | | |

Upper Subansiri Factsheet

Key Indicators

| A) Demographic Indicators (Census 2011) | | | | | | | |
|--|--|----------------|------|----------------|-------|-----------------------------------|------|
| 1 | Population 2011 | 83,205 | | | | | |
| 2 | Sex Ratio | 982 | | | | | |
| B) Infrastructure Details (RHS 2015) | | Numbers | | | | | |
| 3 | Sub Centers (SC) | NA | | | | | |
| 4 | Primary Health Centers (PHC) | NA | | | | | |
| 5 | Community Health Centers (CHC) | 5 | | | | | |
| 6 | SDH | 0 | | | | | |
| 7 | DH | 1 | | | | | |
| C) Maternal Health: | | 2015-16 | | 2016-17 | | | |
| 8 | Expected number of PW | 2,156 | | 2,253 | | | |
| 9 | Expected number of Delivery | 1,999 | | 2,089 | | | |
| | | HMIS (2015-16) | | HMIS (2016-17) | | DLHS-4 (2012-13) | |
| | | Number | PC | Number | PC | | |
| 10 | PW register for ANC | 2,081 | 96.5 | 1,254 | 55.7 | Any ANC | 56.2 |
| 11 | Mothers registered in the first trimester | 570 | 27.4 | 586 | 46.7 | ANC Check up in first trimester | 33.6 |
| 12 | Mothers who had at least 3 Ante-Natal care check ups | 675 | 32.4 | 488 | 38.9 | 3 or more ANC check up | 39.0 |
| 13 | PW who got TT 2 / Booster injection | 719 | 34.6 | 664 | 53.0 | At least 1 TT received | 54.8 |
| 14 | PW who consumed 100 IFA tablets | 602 | 28.9 | 6,279 | 500.7 | 100 IFA tablets consumed | 8.1 |
| 15 | Home Delivery against expected delivery * | 47 | 2.4 | 14 | 0.7 | Full ANC | 6.2 |
| 16 | Institutional Delivery against expected delivery * | 952 | 47.6 | 887 | 42.5 | Home Delivery | 56.2 |
| 17 | Home Delivery against reported delivery ** | 47 | 4.7 | 14 | 1.6 | Institutional Delivery | 43.3 |
| 18 | Institutional Delivery against reported delivery ** | 952 | 95.3 | 887 | 98.4 | Home Delivery by SBA | 0.6 |
| 19 | Total reported Deliveries | 999 | | 901 | | PNC within 14 days of delivery | 37.4 |
| 20 | Reported Delivery at home assisted by a Doctor/Nurse /LHV/ANM(%) | 35 | 74.5 | 11 | 78.6 | | |
| 21 | Woman getting post partum check up between 48 hours and 14 days to total reported deliveries | 884 | 88.5 | 798 | 88.6 | | |
| Note: In case of HMIS data, Percentage is with reference to total number of Pregnant Women registered for ANC during the period and in DLHS-4 is with ref. to Women who had live/still births since three years preceding the survey. (DLHS-4 reference period is from 1-1-2008 to survey date). | | | | | | | |
| 22 | Expected number of Live Birth | 1,960 | | 2,048 | | | |
| D) Child Immunization : | | HMIS (2015-16) | | HMIS (2016-17) | | DLHS-4 (2012-13) | |
| 23 | Total reported live births | 990 | 50.5 | 893 | 43.6 | Breastfeed within 1 hour of birth | 54.4 |
| 24 | Total Still Birth | 16 | 0.8 | 15 | 0.7 | BCG | 77.9 |

| | | | | | | | |
|---|--|------------------------|------|------------------------|------|---------------------------|------|
| 25 | Newborn weighted at birth to live birth | 921 | 93.0 | 873 | 97.8 | 3 dose of DPT | 81.2 |
| 26 | Newborn having weight less than 2.5 kg to newborn weighted at birth. | 15 | 1.6 | 13 | 1.5 | Measles | 63.2 |
| 27 | Newborns breastfed within 1 hour to total reported live birth. | 963 | 97.3 | 877 | 98.2 | Full Immunization | 58 |
| 28 | Fully immunized | 981 | 50.0 | 880 | 43.0 | | |
| 29 | Children who have received BCG | 1,154 | 58.9 | 880 | 43.0 | | |
| 30 | Children who have received DPT1 | 899 | 45.9 | 172 | 8.4 | | |
| 31 | Children who have received DPT3 | 1,039 | 53.0 | 197 | 9.6 | | |
| 32 | Children who have received Measles | 1,082 | 55.2 | 950 | 46.4 | | |
| 33 | Diarrhoea and Dehydration cases reported in children below 5 years | 668 | | 652 | | | |
| <p>Note: In case of HMIS data, Proportions are calculated for 0-11 months old infants to total number of reported live births and in DLHS-4, the percentage of children 12-23 months olds who received specific vaccination to mothers who had last and last but one living child born three years preceding the survey. (DLHS-4 reference period is from 1-1-2008 to survey date).</p> | | | | | | | |
| E) Family Planning | | HMIS (2015-16) | | HMIS (2016-17) | | DLHS-4 (2012-13) | |
| 34 | Number of Vasectomies performed | 0 | | 0 | | Limiting | 8.4 |
| 35 | Number of Tubectomies performed | 0 | | 48 | | Spacing | 23 |
| 36 | Number of IUD insertion | 213 | | 241 | | Women using any method | 47.3 |
| 37 | Number of IUD removals | 212 | | 204 | | Women using modern method | 42.6 |
| F) Service delivery | | HMIS (2015-16) | | HMIS (2016-17) | | | |
| 38 | IPD cases per 10,000 population *** | 177 | | 165 | | | |
| 39 | General OPD cases per 10,000 population *** | 4717 | | 4453 | | | |
| 40 | Ayush OPD cases per 10,000 population *** | 269 | | 295 | | | |
| G) Mortality Details | | | | | | | |
| I) | Infant Deaths (In Numbers) | HMIS (2015-16) | | HMIS (2016-17) | | | |
| 41 | Neonatal death within 24 hours | 0 | | 0 | | | |
| 42 | Up to 1 Weeks of Birth | 2 | | 0 | | | |
| 43 | Between 1 week & 4 weeks of birth | 0 | | 0 | | | |
| 44 | Neonatal Death | 2 | | 0 | | | |
| 45 | Between 1 month and 11 months | 2 | | 0 | | | |
| 46 | Total reported infant deaths | 4 | | 0 | | | |
| II) | Maternal Mortality | HMIS (2015-16) | | HMIS (2016-17) | | | |
| | Maternal Deaths (In Numbers) | HMIS (2015-16) | | HMIS (2016-17) | | | |
| 47 | Due to Abortion | 0 | | 0 | | | |
| 48 | Due to Obstructed/prolonged labour | 0 | | 0 | | | |
| 49 | Due to Severe hypertension/fits | 1 | | 0 | | | |
| 50 | Due to Bleeding | 0 | | 0 | | | |
| 51 | Due to High fever | 0 | | 0 | | | |
| 52 | Due to Other Causes (including causes not known) | 0 | | 0 | | | |
| 53 | Total reported maternal deaths | 1 | | 0 | | | |

West Kameng Factsheet

Key Indicators

| A) Demographic Indicators (Census 2011) | | | | | | | |
|--|--|----------------|-------|----------------|-------|-----------------------------------|------|
| 1 | Population 2011 | 87,013 | | | | | |
| 2 | Sex Ratio | 755 | | | | | |
| B) Infrastructure Details (RHS 2015) | | Numbers | | | | | |
| 3 | Sub Centers (SC) | NA | | | | | |
| 4 | Primary Health Centers (PHC) | NA | | | | | |
| 5 | Community Health Centers (CHC) | 4 | | | | | |
| 6 | SDH | 0 | | | | | |
| 7 | DH | 1 | | | | | |
| C) Maternal Health: | | 2015-16 | | 2016-17 | | | |
| 8 | Expected number of PW | 1,904 | | 1,925 | | | |
| 9 | Expected number of Delivery | 1,765 | | 1,785 | | | |
| | | HMIS (2015-16) | | HMIS (2016-17) | | DLHS-4 (2012-13) | |
| | | Number | PC | Number | PC | | |
| 10 | PW register for ANC | 3,640 | 191.2 | 2,468 | 128.2 | Any ANC | 70.1 |
| 11 | Mothers registered in the first trimester | 1,102 | 30.3 | 998 | 40.4 | ANC Check up in first trimester | 45.7 |
| 12 | Mothers who had at least 3 Ante-Natal care check ups | 1,133 | 31.1 | 1,041 | 42.2 | 3 or more ANC check up | 57.4 |
| 13 | PW who got TT 2 / Booster injection | 1,291 | 35.5 | 1,189 | 48.2 | At least 1 TT received | 68.0 |
| 14 | PW who consumed 100 IFA tablets | 2,001 | 55.0 | 2,321 | 94.0 | 100 IFA tablets consumed | 16.2 |
| 15 | Home Delivery against expected delivery * | 57 | 3.2 | 39 | 2.2 | Full ANC | 14.7 |
| 16 | Institutional Delivery against expected delivery * | 966 | 54.7 | 943 | 52.8 | Home Delivery | 50.4 |
| 17 | Home Delivery against reported delivery ** | 57 | 5.6 | 39 | 4.0 | Institutional Delivery | 49.3 |
| 18 | Institutional Delivery against reported delivery ** | 966 | 94.4 | 943 | 96.0 | Home Delivery by SBA | 2.4 |
| 19 | Total reported Deliveries | 1,023 | | 982 | | PNC within 14 days of delivery | 64.1 |
| 20 | Reported Delivery at home assisted by a Doctor/Nurse /LHV/ANM(%) | 52 | 91.2 | 30 | 76.9 | | |
| 21 | Woman getting post partum check up between 48 hours and 14 days to total reported deliveries | 532 | 52.0 | 621 | 63.2 | | |
| Note: In case of HMIS data, Percentage is with reference to total number of Pregnant Women registered for ANC during the period and in DLHS-4 is with ref. to Women who had live/still births since three years preceding the survey. (DLHS-4 reference period is from 1-1-2008 to survey date). | | | | | | | |
| 22 | Expected number of Live Birth | 1,731 | | 1,750 | | | |
| D) Child Immunization : | | HMIS (2015-16) | | HMIS (2016-17) | | DLHS-4 (2012-13) | |
| 23 | Total reported live births | 975 | 56.3 | 983 | 56.2 | Breastfeed within 1 hour of birth | 56.7 |
| 24 | Total Still Birth | 22 | 1.3 | 7 | 0.4 | BCG | 86.7 |

| | | | | | | | |
|---|--|------------------------|------|------------------------|------|---------------------------|------|
| 25 | Newborn weighted at birth to live birth | 926 | 95.0 | 954 | 97.0 | 3 dose of DPT | 85.3 |
| 26 | Newborn having weight less than 2.5 kg to newborn weighted at birth. | 39 | 4.2 | 41 | 4.3 | Measles | 75 |
| 27 | Newborns breastfed within 1 hour to total reported live birth. | 940 | 96.4 | 952 | 96.8 | Full Immunization | 61.8 |
| 28 | Fully immunized | 1,263 | 73.0 | 1,264 | 72.2 | | |
| 29 | Children who have received BCG | 1,292 | 74.6 | 1,036 | 59.2 | | |
| 30 | Children who have received DPT1 | 1,326 | 76.6 | 51 | 2.9 | | |
| 31 | Children who have received DPT3 | 1,187 | 68.6 | 71 | 4.1 | | |
| 32 | Children who have received Measles | 1,208 | 69.8 | 1,332 | 76.1 | | |
| 33 | Diarrhoea and Dehydration cases reported in children below 5 years | 1,082 | | 840 | | | |
| <p>Note: In case of HMIS data, Proportions are calculated for 0-11 months old infants to total number of reported live births and in DLHS-4, the percentage of children 12-23 months olds who received specific vaccination to mothers who had last and last but one living child born three years preceding the survey. (DLHS-4 reference period is from 1-1-2008 to survey date).</p> | | | | | | | |
| E) Family Planning | | HMIS (2015-16) | | HMIS (2016-17) | | DLHS-4 (2012-13) | |
| 34 | Number of Vasectomies performed | 0 | | 0 | | Limiting | 11.5 |
| 35 | Number of Tubectomies performed | 15 | | 16 | | Spacing | 15.8 |
| 36 | Number of IUD insertion | 554 | | 535 | | Women using any method | 55.4 |
| 37 | Number of IUD removals | 227 | | 267 | | Women using modern method | 49 |
| F) Service delivery | | HMIS (2015-16) | | HMIS (2016-17) | | | |
| 38 | IPD cases per 10,000 population *** | 325 | | 313 | | | |
| 39 | General OPD cases per 10,000 population *** | 8410 | | 9204 | | | |
| 40 | Ayush OPD cases per 10,000 population *** | 131 | | 174 | | | |
| G) Mortality Details | | | | | | | |
| I) | Infant Deaths (In Numbers) | HMIS (2015-16) | | HMIS (2016-17) | | | |
| 41 | Neonatal death within 24 hours | 0 | | 0 | | | |
| 42 | Up to 1 Weeks of Birth | 0 | | 0 | | | |
| 43 | Between 1 week & 4 weeks of birth | 0 | | 0 | | | |
| 44 | Neonatal Death | 0 | | 0 | | | |
| 45 | Between 1 month and 11 months | 0 | | 0 | | | |
| 46 | Total reported infant deaths | 0 | | 0 | | | |
| II) | Maternal Mortality | | | | | | |
| | Maternal Deaths (In Numbers) | HMIS (2015-16) | | HMIS (2016-17) | | | |
| 47 | Due to Abortion | 0 | | 0 | | | |
| 48 | Due to Obstructed/prolonged labour | 0 | | 0 | | | |
| 49 | Due to Severe hypertension/fits | 0 | | 0 | | | |
| 50 | Due to Bleeding | 0 | | 0 | | | |
| 51 | Due to High fever | 0 | | 0 | | | |
| 52 | Due to Other Causes (including causes not known) | 0 | | 0 | | | |
| 53 | Total reported maternal deaths | 0 | | 0 | | | |

West Siang Factsheet

Key Indicators

| A) Demographic Indicators (Census 2011) | | | | | | | |
|--|--|-----------------|------|-----------------|-------|-----------------------------------|------|
| 1 | Population 2011 | 112,272 | | | | | |
| 2 | Sex Ratio | 916 | | | | | |
| B) Infrastructure Details (RHS 2015) | | Numbers | | | | | |
| 3 | Sub Centers (SC) | NA | | | | | |
| 4 | Primary Health Centers (PHC) | NA | | | | | |
| 5 | Community Health Centers (CHC) | 6 | | | | | |
| 6 | SDH | 0 | | | | | |
| 7 | DH | 1 | | | | | |
| C) Maternal Health: | | 2015-16 | | 2016-17 | | | |
| 8 | Expected number of PW | 2,461 | | 2,468 | | | |
| 9 | Expected number of Delivery | 2,282 | | 2,289 | | | |
| | | HMIS (2015-16) | | HMIS (2016-17) | | DLHS-4 (2012-13) | |
| | | Number | PC | Number | PC | | |
| 10 | PW register for ANC | 1,930 | 78.4 | 2,053 | 83.2 | Any ANC | 71.5 |
| 11 | Mothers registered in the first trimester | 448 | 23.2 | 519 | 25.3 | ANC Check up in first trimester | 31.9 |
| 12 | Mothers who had at least 3 Ante-Natal care check ups | 953 | 49.4 | 1,003 | 48.9 | 3 or more ANC check up | 58.6 |
| 13 | PW who got TT 2 / Booster injection | 1,102 | 57.1 | 1,019 | 49.6 | At least 1 TT received | 65.6 |
| 14 | PW who consumed 100 IFA tablets | 1,289 | 66.8 | 1,468 | 71.5 | 100 IFA tablets consumed | 30.6 |
| 15 | Home Delivery against expected delivery * | 5 | 0.2 | 3 | 0.1 | Full ANC | 27.0 |
| 16 | Institutional Delivery against expected delivery * | 1,247 | 54.6 | 1,281 | 56.0 | Home Delivery | 40.7 |
| 17 | Home Delivery against reported delivery ** | 5 | 0.4 | 3 | 0.2 | Institutional Delivery | 58.6 |
| 18 | Institutional Delivery against reported delivery ** | 1,247 | 99.6 | 1,281 | 99.8 | Home Delivery by SBA | 0.7 |
| 19 | Total reported Deliveries | 1,252 | | 1,284 | | PNC within 14 days of delivery | 31.5 |
| 20 | Reported Delivery at home assisted by a Doctor/Nurse /LHV/ANM(%) | 2 | 40.0 | 3 | 100.0 | | |
| 21 | Woman getting post partum check up between 48 hours and 14 days to total reported deliveries | 341 | 27.2 | 159 | 12.4 | | |
| Note: In case of HMIS data, Percentage is with reference to total number of Pregnant Women registered for ANC during the period and in DLHS-4 is with ref. to Women who had live/still births since three years preceding the survey. (DLHS-4 reference period is from 1-1-2008 to survey date). | | | | | | | |
| 22 | Expected number of Live Birth | 2,237 | | 2,244 | | | |
| D) Child Immunization : | | HMIS (2015-16) | | HMIS (2016-17) | | DLHS-4 (2012-13) | |
| 23 | Total reported live births | 1,238 | 55.3 | 1,266 | 56.4 | Breastfeed within 1 hour of birth | 59.7 |
| 24 | Total Still Birth | 15 | 0.7 | 17 | 0.8 | BCG | 68.9 |

| | | | | | | | |
|---|--|------------------------|------|------------------------|------|---------------------------|------|
| 25 | Newborn weighted at birth to live birth | 1,223 | 98.8 | 1,225 | 96.8 | 3 dose of DPT | 58.9 |
| 26 | Newborn having weight less than 2.5 kg to newborn weighted at birth. | 46 | 3.8 | 49 | 4.0 | Measles | 66.2 |
| 27 | Newborns breastfed within 1 hour to total reported live birth. | 1,156 | 93.4 | 1,149 | 90.8 | Full Immunization | 50 |
| 28 | Fully immunized | 1,305 | 58.3 | 1,250 | 55.7 | | |
| 29 | Children who have received BCG | 1,500 | 67.0 | 1,324 | 59.0 | | |
| 30 | Children who have received DPT1 | 1,050 | 46.9 | 18 | 0.8 | | |
| 31 | Children who have received DPT3 | 1,240 | 55.4 | 15 | 0.7 | | |
| 32 | Children who have received Measles | 1,385 | 61.9 | 1,367 | 60.9 | | |
| 33 | Diarrhoea and Dehydration cases reported in children below 5 years | 686 | | 942 | | | |
| <p>Note: In case of HMIS data, Proportions are calculated for 0-11 months old infants to total number of reported live births and in DLHS-4, the percentage of children 12-23 months olds who received specific vaccination to mothers who had last and last but one living child born three years preceding the survey. (DLHS-4 reference period is from 1-1-2008 to survey date).</p> | | | | | | | |
| E) Family Planning | | HMIS (2015-16) | | HMIS (2016-17) | | DLHS-4 (2012-13) | |
| 34 | Number of Vasectomies performed | 0 | | 0 | | Limiting | 11.1 |
| 35 | Number of Tubectomies performed | 89 | | 58 | | Spacing | 13.6 |
| 36 | Number of IUD insertion | 344 | | 291 | | Women using any method | 61.3 |
| 37 | Number of IUD removals | 192 | | 218 | | Women using modern method | 55.2 |
| F) Service delivery | | HMIS (2015-16) | | HMIS (2016-17) | | | |
| 38 | IPD cases per 10,000 population *** | 201 | | 226 | | | |
| 39 | General OPD cases per 10,000 population *** | 7443 | | 7641 | | | |
| 40 | Ayush OPD cases per 10,000 population *** | 363 | | 359 | | | |
| G) Mortality Details | | | | | | | |
| I) | Infant Deaths (In Numbers) | HMIS (2015-16) | | HMIS (2016-17) | | | |
| 41 | Neonatal death within 24 hours | 0 | | 0 | | | |
| 42 | Up to 1 Weeks of Birth | 0 | | 0 | | | |
| 43 | Between 1 week & 4 weeks of birth | 0 | | 0 | | | |
| 44 | Neonatal Death | 0 | | 0 | | | |
| 45 | Between 1 month and 11 months | 0 | | 0 | | | |
| 46 | Total reported infant deaths | 0 | | 0 | | | |
| II) | Maternal Mortality | HMIS (2015-16) | | HMIS (2016-17) | | | |
| | Maternal Deaths (In Numbers) | HMIS (2015-16) | | HMIS (2016-17) | | | |
| 47 | Due to Abortion | 0 | | 0 | | | |
| 48 | Due to Obstructed/prolonged labour | 0 | | 0 | | | |
| 49 | Due to Severe hypertension/fits | 0 | | 0 | | | |
| 50 | Due to Bleeding | 0 | | 0 | | | |
| 51 | Due to High fever | 0 | | 0 | | | |
| 52 | Due to Other Causes (including causes not known) | 0 | | 0 | | | |
| 53 | Total reported maternal deaths | 0 | | 0 | | | |